

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|----|--|
| (51) International Patent Classification ⁶ : C12Q 1/68, 1/70 | A1 | (11) International Publication Number: WO 97/21836 (43) International Publication Date: 19 June 1997 (19.06.97) |
| (21) International Application Number: PCT/US96/19680 (22) International Filing Date: 13 December 1996 (13.12.96) (30) Priority Data: 08/572,645 14 December 1995 (14.12.95) US (71) Applicant: THE GENERAL HOSPITAL CORPORATION [US/US]; 55 Fruit Street, Boston, MA 02110 (US). (72) Inventor: FISHMAN, Jay, A.; 383 Linden Street, Wellesley, MA 02181 (US). (74) Agents: MYERS, Paul, Louis et al.; Lahive & Cockfield L.L.P., 60 State Street, Boston, MA 02109 (US). | | (81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> |
| (54) Title: MOLECULAR SEQUENCE OF SWINE RETROVIRUS AND METHODS OF USE (57) Abstract <p>Purified nucleic acid which can specifically hybridize with the sequence of swine retroviruses.</p> | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|--|----|--------------------------|
| AM | Armenia | GB | United Kingdom | MW | Malawi |
| AT | Austria | GE | Georgia | MX | Mexico |
| AU | Australia | GN | Guinea | NE | Niger |
| BB | Barbados | GR | Greece | NL | Netherlands |
| BE | Belgium | HU | Hungary | NO | Norway |
| BF | Burkina Faso | IE | Ireland | NZ | New Zealand |
| BG | Bulgaria | IT | Italy | PL | Poland |
| BJ | Benin | JP | Japan | PT | Portugal |
| BR | Brazil | KE | Kenya | RO | Romania |
| BY | Belarus | KG | Kyrgyzstan | RU | Russian Federation |
| CA | Canada | KP | Democratic People's Republic of Korea | SD | Sudan |
| CF | Central African Republic | KR | Republic of Korea | SE | Sweden |
| CG | Congo | KZ | Kazakhstan | SG | Singapore |
| CH | Switzerland | LI | Liechtenstein | SI | Slovenia |
| CI | Côte d'Ivoire | LK | Sri Lanka | SK | Slovakia |
| CM | Cameroon | LR | Liberia | SN | Senegal |
| CN | China | LT | Lithuania | SZ | Swaziland |
| CS | Czechoslovakia | LU | Luxembourg | TD | Chad |
| CZ | Czech Republic | LV | Latvia | TG | Togo |
| DE | Germany | MC | Monaco | TJ | Tajikistan |
| DK | Denmark | MD | Republic of Moldova | TT | Trinidad and Tobago |
| EE | Estonia | MG | Madagascar | UA | Ukraine |
| ES | Spain | ML | Mali | UG | Uganda |
| FI | Finland | MN | Mongolia | US | United States of America |
| FR | France | MR | Mauritania | UZ | Uzbekistan |
| GA | Gabon | | | VN | Viet Nam |

MOLECULAR SEQUENCE OF SWINE RETROVIRUS AND METHODS OF USE

5 This application is a continuation-in-part of U.S.S.N. 08/572,645, filed December 14, 1995, which is hereby incorporated by reference.

Field of the Invention

 The invention relates to porcine retroviral sequences, peptides encoded by porcine retroviral sequences, and methods of using the porcine retroviral nucleic acids and peptides.

10 Background of the Invention

 Advances in solid organ transplantation and a chronic shortage of suitable organ donors have made xenotransplantation an attractive alternative to the use of human allografts. However, the potential for introduction of a new group of infectious diseases from donor animals into the human population is a concern with the use of these methods.

15 The term applied to the natural acquisition by humans of infectious agents carried by other species is zoonosis. The transplantation of infection from nonhuman species into humans is best termed "direct zoonosis" or "xenosis."

 Nonhuman primates and swine have been considered the main potential sources of organs for xenotransplantation (Niekrasz et al. (1992) *Transplant Proc* 24:625; Starzl et al. (1993) *Lancet* 341:65; Murphy et al. (1970) *Trans Proc* 4:546; Brede and Murphy (1972) *Primates Med* 7:18; Cooper et al. In Xenotransplantation: The Transplantation of Organs and Tissues between Species, eds. Cooper et al. (1991) p. 457; RY Calne (1970) *Transplant Proc* 2:550; H. Auchincloss, Jr. (1988) *Transplantation* 46:1; and Chiche et al. (1993) *Transplantation* 6:1418). The infectious disease issues for primates and swine are similar to those of human donors. The prevention of infection depends on the ability to predict, to recognize, and to prevent common infections in the immunocompromised transplantation recipient (Rubin et al. (1993) *Antimicrob Agents Chemother* 37:619). Because of the potential carriage by nonhuman primates of pathogens easily adopted to humans, ethical concerns, and the cost of maintaining large colonies of primates, other species have received consideration as organ donors (Brede and Murphy (1972) *Primates Med* 7:18; Van Der Riet et al. (1987) *Transplant Proc* 19:4069; Katler In Xenotransplantation: The Transplantation of Organs and Tissues between Species, eds. Cooper et al. (1991) p. 457; Metzger et al. (1981) *J Immunol* 127:769; McClure et al. (1987) *Nature* 330:487; Letvin et al. (1987) *J Infect Dis* 156:406; Castro et al. (1991) *Virology* 184:219; Benveniste and Todaro (1973) *Proc Natl Acad Sci USA* 70:3316; and Teich, in RNA Tumor viruses, eds. Weiss et. al. (1985) p. 25) The economic importance of swine and experience in studies of transplantation in the miniature swine model have allowed some of the potential pathogens associated with these animals to be defined (Niekrasz et al. (1992) *Transplant Proc* 24:625;

Cooper et al. In Xenotransplantation: The Transplantation of Organs and Tissues between Species, eds. Cooper et al. (1991) p. 457; and Leman et al. (1992) Diseases of Swine. 7th ed. Ames, Iowa: Iowa State University). Miniature swine have received consideration as organ donors because of a number of features of the species. The structure and function of the main pig organs are comparable to those of man. Swine attain body weights and organ sizes adequate to the provision of organs for human use. Lastly, veterinarians and commercial breeders have developed approaches to creation of specific-pathogen-free (SPF) swine with the ability to eliminate known pathogens from breeding colonies (Alexander et al. (1980) *Proc 6th Int Congr Pig Vet Soc*, Copenhagen; Betts (1961) *Vet Rec* 73:1349; Betts et al. (1960) *Vet Rec* 72:461; Caldwell et al. (1959) *J Am Vet Med Assoc* 135:504; and Yong (1964) *Adv Vet Sci* 9:61).

Concern exists over the transfer of porcine retroviruses by xenotransplantation (Smith (1993) *N Engl J Med* 328:141). Many of the unique properties of the retroviruses are due to the synthesis of a complementary DNA copy from the RNA template (by reverse transcriptase), and integration of this DNA into the host genome. The integrated retroviral copy (which is referred to as an endogenous copy or "provirus") can be transmitted via the germ line.

Summary of the Invention

In general, the invention features a purified swine or miniature swine retroviral nucleic acid, e.g., a Tsukuba nucleic acid, a purified miniature swine retroviral nucleic acid sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, and methods of their use in detecting the presence of porcine, e.g., miniature swine, retroviral sequences.

In another aspect, the invention features a purified nucleic acid, e.g., a probe or primer, which can specifically hybridize with a purified swine or miniature swine retroviral genome, e.g., a Tsukuba genome, the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In preferred embodiments the nucleic acid is other than the entire retroviral genome of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, e.g., it is at least 1 nucleotide longer, or at least 1 nucleotide shorter, or differs in sequence at at least one position, e.g., the nucleic acid is a fragment of the sequence of SEQ ID NO:1 or its complement SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, or it includes sequence additional to that of SEQ ID NO:1, or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In preferred embodiments, the nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from

SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In other embodiments: the sequence of the nucleic acid differs from the corresponding sequence of SEQ ID NO: 1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, by 1, 2, 3, 4, or 5 base pairs; the sequence of the nucleic acid differs from the corresponding sequence of SEQ ID NO: 1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, by at least 1, 2, 3, 4, or 5 base pairs but less than 6, 7, 8, 9, or 10 base pairs.

In other preferred embodiments: the nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length.

In yet other preferred embodiments: the nucleic acid can specifically hybridize with a translatable region of a miniature swine retroviral genome, e.g., the retroviral genome of SEQ ID NO: 1, or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, e.g., a region from the gag, pol, or env gene; the probe or primer can specifically hybridize with an untranslated region of a miniature swine retroviral genome, e.g., the retroviral genome of SEQ ID NO: 1, or its complement SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement; the probe or primer can specifically hybridize with a non-conserved region of a miniature swine retroviral genome, e.g., the retroviral genome of SEQ ID NO: 1, or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement; the probe or primer can specifically hybridize with the highly conserved regions of a miniature swine retroviral genome, e.g., the retroviral genome of SEQ ID NO: 1, or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In preferred embodiments, the primer is selected from the group consisting of SEQ ID NOs:4-74.

In preferred embodiments, hybridization of the probe to retroviral sequences can be detected by standard methods, e.g., by radiolabeled probes or by probes bearing nonradioactive markers such as enzymes or antibody binding sites. For example, a probe can be conjugated with an enzyme such as horseradish peroxidase, where the enzymatic activity of the conjugated enzyme is used as a signal for hybridization. Alternatively, the probe can be coupled to an epitope recognized by an antibody, e.g., an antibody conjugated to an enzyme or another marker.

In another aspect, the invention features a reaction mixture which includes a target nucleic acid, e.g., a human, swine, or a miniature swine nucleic acid, and a purified second nucleic acid, e.g., a probe or primer, as, e.g., is described herein, which specifically

hybridizes with the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, a swine or a miniature swine retroviral nucleic acid, e.g., a Tsukuba nucleic acid.

In preferred embodiments, the target nucleic acid: includes RNA; or includes DNA.

5 In preferred embodiments, the target nucleic acid includes: genomic DNA isolated from a miniature swine; RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine; DNA, RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine organ, e.g., a kidney; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine potential donor organ; RNA,
10 DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine organ which has been transplanted into a organ recipient, e.g., a xenogeneic recipient, e.g., a primate, e.g., a human.

In preferred embodiments, the target nucleic acid includes: genomic DNA isolated from a swine; RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a
15 swine; DNA, RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine organ, e.g., a kidney; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine potential donor organ; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine organ which has been transplanted into a organ recipient, e.g., a xenogeneic recipient, e.g., a primate, e.g., a human.

20 In a preferred embodiment: the second nucleic acid is a porcine retroviral sequence, probe or primer, e.g., as described herein, e.g., a Tsukuba-1 retroviral sequence; the second nucleic acid is a sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, or a fragment of the sequence or complement at least 10, 20, or 30, basepairs in length.

25 In preferred embodiments, the second nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

30 In other preferred embodiments: the second nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the second nucleic acid is a full length retroviral
35 genome.

In preferred embodiments the second nucleic acid is: a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from

nucleotides 2452-4839 (e.g. from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g. from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g. from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g. from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g. from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof.

In another aspect, the invention features a method for screening a cell or a tissue, e.g., a cellular or tissue transplant, e.g., a xenograft, for the presence or expression of a swine or a miniature swine retrovirus or retroviral sequence, e.g., an endogenous miniature swine retrovirus. The method includes:

contacting a target nucleic acid from the tissue with a second sequence chosen from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g. from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g. from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g. from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g. from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g. from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid under conditions in which hybridization can occur, hybridization being indicative of the presence or expression of an

endogenous miniature swine retrovirus or retroviral sequence in the tissue or an endogenous swine retrovirus in the tissue.

In preferred embodiments, the method further includes amplifying the target nucleic acid with primers which specifically hybridize to the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In preferred embodiments, the tissue or cellular transplant is selected from the group consisting of: heart, lung, liver, bone marrow, kidney, brain cells, neural tissue, pancreas or pancreatic cells, thymus, or intestinal tissue.

In other preferred embodiments, the target nucleic acid is: DNA; RNA; or cDNA.

In other preferred embodiments, the target nucleic acid is taken from: a tissue sample, or a blood sample, e.g., a tissue biopsy sample, e.g., a tissue sample suitable for *in situ* hybridization or immunohistochemistry.

In preferred embodiments, the target nucleic acid includes: genomic DNA isolated from a miniature swine; RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine; DNA, RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine organ, e.g., a kidney; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine potential donor organ; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine organ which has been transplanted into a organ recipient, e.g., a xenogeneic recipient, e.g., a primate, e.g., a human.

In preferred embodiments, the target nucleic acid includes: genomic DNA isolated from a swine; RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine; DNA, RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine organ, e.g., a kidney; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine potential donor organ; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine organ which has been transplanted into a organ recipient, e.g., a recipient swine or a xenogeneic recipient, e.g., a primate, e.g., a human.

In a preferred embodiment the target nucleic acid is RNA, or a nucleic acid amplified from RNA in the tissue, and hybridization is correlated with expression of an endogenous miniature swine retrovirus or retroviral sequence or an endogenous swine retrovirus.

In a preferred embodiment the target nucleic acid is DNA, or a nucleic acid amplified from DNA in the tissue, and hybridization is correlated with the presence of an endogenous miniature swine retrovirus or an endogenous swine retrovirus.

In preferred embodiments, the second nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence

from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In other preferred embodiments: the second nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the second nucleic acid is a full length retroviral genome.

In another aspect, the invention features a method of screening a porcine derived cell or tissue for the presence of an activatable porcine retrovirus, e.g., an activatable porcine provirus. The method includes:

stimulating a porcine derived cell or tissue with a treatment which can activate a retrovirus;

contacting a target nucleic acid from the porcine derived cell or tissue with a second sequence chosen from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g, from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g, from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid hybridization being indicative of the presence of an activatable porcine provirus in the porcine derived cell or tissue.

In preferred embodiments the treatment is: contact with a drug, e.g., a steroid or a cytotoxic agent, infection or contact with a virus, the induction of stress, e.g., nutritional stress or immunologic stress, e.g., contact with a T-cell, e.g., a reactive T-cell.

5 In preferred embodiments, the method further includes amplifying the target nucleic acid with primers which specifically hybridize to the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In other preferred embodiments, the target nucleic acid is taken from: a tissue sample, or a blood sample, e.g., a tissue biopsy sample, e.g., a tissue sample suitable for *in situ* hybridization or immunohistochemistry.

10 In preferred embodiments, the target nucleic acid includes: genomic DNA isolated from a miniature swine; RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine; DNA, RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine organ, e.g., a kidney; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine potential donor organ; RNA,
15 DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine organ which has been transplanted into a organ recipient, e.g., a xenogeneic recipient, e.g., a primate, e.g., a human.

In preferred embodiments, the target nucleic acid includes: genomic DNA isolated from a swine; RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a
20 swine; DNA, RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine organ, e.g., a kidney; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine potential donor organ; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine organ which has been transplanted into a organ recipient, e.g., a recipient swine or a xenogeneic recipient, e.g., a primate, e.g., a
25 human.

In preferred embodiments, the second nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3
30 or its complement.

In other preferred embodiments: the second nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000,
35 6000, or 8060 nucleotides in length; the second nucleic acid is a full length retroviral genome.

In another aspect, the invention features a method for screening a miniature swine genome or a swine genome for the presence of a porcine retrovirus or retroviral sequence, e.g., an endogenous porcine retrovirus. The method includes:

contacting the miniature swine (or swine) genomic DNA with a second sequence
5 chosen from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the
10 sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g. from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g. from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g. from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at
15 least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a
20 env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g. from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g. from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid under conditions in which the
25 sequences can hybridize, hybridization being indicative of the presence of the endogenous porcine retrovirus or retroviral sequence in the miniature swine (or swine) genome.

In preferred embodiments, the method further includes amplifying all or a portion of the miniature swine (or swine) genome with primers which specifically hybridize to the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID
30 NO:3 or its complement.

In a preferred embodiment: the second nucleic acid is a porcine retroviral sequence, probe or primer, e.g., as described herein, e.g., a Tsukuba-1 retroviral sequence; the second nucleic acid is a sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, or a fragment of the sequence or
35 complement at least 10, 20, or 30, basepairs in length.

In preferred embodiments, the second nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence

from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In other preferred embodiments: the second nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the second nucleic acid is a full length retroviral genome.

In another aspect, the invention features a method for screening a genetically modified miniature swine or a genetically modified swine for the presence or expression of a miniature swine or swine retrovirus or retroviral sequence, e.g., an endogenous miniature swine retrovirus. The method includes:

contacting a target nucleic acid from the genetically modified miniature swine or swine with a second sequence chosen from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g, from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g, from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof;

a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid under conditions in which hybridization can occur, hybridization being indicative of the presence or expression of an endogenous miniature swine retrovirus or retroviral sequence or swine retrovirus or retroviral sequence in the genetically modified miniature swine or swine.

In preferred embodiments, the method further includes amplifying the target nucleic acid with primers which specifically hybridize to the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

5 In preferred embodiments, the target nucleic acid includes: genomic DNA isolated from a miniature swine; RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine; DNA, RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine organ, e.g., a kidney; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine potential donor organ; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine organ which has been transplanted into a organ recipient, e.g., a xenogeneic recipient, e.g., a primate, e.g., a human.

10 In preferred embodiments, the target nucleic acid includes: genomic DNA isolated from a swine; RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine; DNA, RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine organ, e.g., a kidney; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine potential donor organ; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine organ which has been transplanted into a organ recipient, e.g., a recipient swine or a xenogeneic recipient, e.g., a primate, e.g., a human.

20 In preferred embodiments, the second nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

25 In other preferred embodiments: the second nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the second nucleic acid is a full length retroviral genome.

30 In another aspect, the invention features a method of assessing the potential risk associated with the transplantation of a graft from a donor miniature swine or swine into a recipient animal, e.g., a miniature swine or swine, a non-human primate, or a human. The method includes:

35 contacting a target nucleic acid from the donor, recipient or the graft, with a second sequence chosen from the group of: a nucleic acid sequence which specifically hybridizes a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a

sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive
5 nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g. from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g. from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g. from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof;

a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which
10 encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof;

a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which
15 encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g. from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g. from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid under conditions in
20 which the sequences can hybridize, hybridization being indicative of a risk associated with the transplantation.

In a preferred embodiment: the second nucleic acid is a Tsukuba-1 retroviral sequence, probe or primer, e.g., as described herein; the second nucleic acid is a porcine retroviral sequence, probe or primer, e.g., as described herein; the second nucleic acid is the
25 sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, or a fragment of the sequence or complement at least 10, 20, or 30, basepairs in length.

In preferred embodiments, the target nucleic acid includes: genomic DNA isolated from a miniature swine; RNA or cDNA, e.g., cDNA made from an RNA template, isolated
30 from a miniature swine; DNA, RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine organ, e.g., a kidney; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine potential donor organ; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine organ which has been transplanted into a organ recipient, e.g., a xenogeneic recipient, e.g., a
35 primate, e.g., a human.

In preferred embodiments, the target nucleic acid includes: genomic DNA isolated from a swine; RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine; DNA, RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a

swine organ, e.g., a kidney; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine potential donor organ; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine organ which has been transplanted into a organ recipient, e.g., a recipient swine or a xenogeneic recipient, e.g., a primate, e.g., a human.

In preferred embodiments, the second nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In other preferred embodiments: the second nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the second nucleic acid is a full length retroviral genome.

In another aspect, the invention features a method of determining if an endogenous miniature swine or swine retrovirus or retroviral sequence genome includes a mutation which modulates its expression, e.g., results in misexpression. The method includes: determining the structure of the endogenous retroviral genome, and comparing the structure of the endogenous retroviral genome with the retroviral sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, a difference being predictive of a mutation.

In preferred embodiments the method includes sequencing the endogenous genome and comparing it with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In preferred embodiments, the method includes using primers to amplify, e.g., by PCR, LCR (ligase chain reaction), or other amplification methods, a region of the endogenous retroviral genome, and comparing the structure of the amplification product to the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement to determine if there is difference in sequence between retroviral genome and SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement. The method further includes determining if one or more restriction sites exist in the endogenous retroviral genome, and determining if the sites exist in SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In preferred embodiments, the mutation is a gross defect, e.g., an insertion, inversion, translocation or a deletion, of all or part of the retroviral genome.

In preferred embodiments, detecting the mutation can include: (i) providing a labeled PCR probe amplified from DNA (e.g., SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3) containing a porcine retroviral nucleotide sequence which hybridizes to a sense or antisense sequence from the porcine retroviral genome (e.g., SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3), or naturally occurring mutants thereof; (ii) exposing the probe/primer to nucleic acid of the tissue (e.g., genomic DNA) digested with a restriction endonuclease; and (iii) detecting by *in situ* hybridization of the probe/primer to the nucleic acid, the presence or absence of the genetic lesion. Alternatively, direct PCR analysis, using primers specific for porcine retroviral genes (e.g., genes comprising the nucleotide sequence shown in SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3), can be used to detect the presence or absence of the genetic lesion in the porcine retroviral genome by comparing the products amplified.

In another aspect, the invention features a method of providing a miniature swine or a swine free of an endogenous retrovirus or retroviral sequence, e.g., activatable retrovirus, insertion at a preselected site. The method includes:

- 15 performing a breeding cross between a first miniature swine (or swine) having a retroviral insertion at the preselected site and a second miniature swine (or swine) not having a retroviral insertion at a preselected site, e.g., the same site, and recovering a progeny miniature swine (or swine), not having the insertion, wherein the presence or absence of the retroviral insertion is determined by contacting the genome of a miniature swine (or swine) with a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g., from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g., from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g., from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof;
- 25 a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g., from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g., from nucleotides 4738-6722) of SEQ ID NO:2.
- 30
- 35

or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof: a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid.

In preferred embodiments, the nucleic acid is hybridized to nucleic acid, e.g., DNA from the genome, of the first animal or one of its ancestors.

5 In preferred embodiments, the nucleic acid is hybridized to nucleic acid, e.g., DNA from the genome, of the second animal or one of its ancestors.

In preferred embodiments, the nucleic acid is hybridized to nucleic acid, e.g., DNA from the genome, of the progeny animal or one of its descendants.

10 In preferred embodiments, the nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

15 In other preferred embodiments: the nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is a full length retroviral genome.

20 In another aspect, the invention features a method of evaluating a treatment, e.g., an immunosuppressive treatment, for the ability to activate a retrovirus, e.g., an endogenous porcine retrovirus. The method includes:

administering a treatment to a subject, e.g., a miniature swine (or a swine), having an endogenous porcine retrovirus; and

25 detecting expression of the porcine retrovirus with a purified nucleic acid sequence which specifically hybridizes to the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In preferred embodiments, the immunosuppressive treatment includes radiation, chemotherapy or drug treatment.

30 In preferred embodiments: the treatment is one which can induce immunological tolerance; the treatment is one which can introduce new genetic material, e.g., introduce new genetic material into a miniature swine genome (or a swine genome) or into the genome of a host which receives a swine or a miniature swine graft, e.g., the treatment is one which introduces a new genetic material via retroviral mediated transfer.

35 In a preferred embodiment: the purified nucleic acid is a Tsukuba-1 retroviral sequence, probe or primer, e.g., as described herein; the purified nucleic acid is a porcine retroviral sequence, probe or primer, e.g., as described herein; the purified nucleic acid is the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ

ID NO:3 or its complement, or a fragment of such sequence or complement at least 10, 20, or 30, basepairs in length.

In preferred embodiments, the purified nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%,
5 most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In other preferred embodiments: the purified nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100,
10 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the purified nucleic acid is a full length retroviral genome.

In preferred embodiments the second nucleic acid is: a nucleic acid of at least 10
15 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g, from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g, from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof;
20 a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense
25 sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof.

In another aspect, the invention features a method of localizing the origin of a
30 porcine retroviral infection. The method includes:

contacting a target nucleic acid from the graft with a second sequence chosen from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2
35 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g, from

nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g, from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a

5 nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof;

a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or

10 antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid contacting a target nucleic acid from the recipient with a second sequence chosen from the group of: a

15 sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or

20 antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g, from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g, from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10

25 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a

30 env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid; hybridization to the nucleic acid

35 from the graft correlates with the porcine retroviral infection in the graft; and hybridization to the nucleic acid from the recipient correlates with the porcine retroviral infection in the recipient.

In preferred embodiments, the target nucleic acid includes: genomic DNA, RNA or cDNA, e.g., cDNA made from an RNA template.

In a preferred embodiment: the second nucleic acid is a porcine retroviral sequence, probe or primer, e.g., as described herein, e.g., a Tsukuba-1 retroviral sequence; the second
5 nucleic acid is a sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, or a fragment of the sequence or complement at least 10, 20, or 30, basepairs in length.

In preferred embodiments, the recipient is an animal, e.g., a miniature swine, a swine, a non-human primate, or a human.

10 In preferred embodiments, the graft is selected from the group consisting of: heart, lung, liver, bone marrow or kidney.

In preferred embodiments, the second nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most
15 preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In other preferred embodiments: the second nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100,
20 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the second nucleic acid is a full length retroviral genome.

In another aspect, the invention features a method of screening a cell, e.g., a cell
25 having a disorder, e.g., a proliferative disorder, e.g., a tumor cell, e.g., a cancer cell, e.g., a lymphoma or a hepatocellular carcinoma, developing in a graft recipient, e.g., a xenograft, for the presence or expression of a porcine retrovirus or retroviral sequence. The method includes:

contacting a target nucleic acid from a tumor cell with a second sequence chosen
30 from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of
35 sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g., from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g., from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g., from nucleotides

585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or
5 nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides
10 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid, under conditions in which the sample and the nucleic acid sequence can hybridize, hybridization being indicative of the presence of the endogenous porcine retrovirus or retroviral sequence in the tumor cell.

In preferred embodiments, the target nucleic acid from a tumor cell includes:
15 genomic DNA, RNA or cDNA, e.g., cDNA made from an RNA template.

In a preferred embodiment: the second nucleic acid is a porcine retroviral sequence, probe or primer, e.g., as described herein, e.g., a Tsukuba-1 retroviral sequence; the second nucleic acid is a sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, or a fragment of the sequence or
20 complement at least 10, 20, or 30, basepairs in length.

In preferred embodiments, the second nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3
25 or its complement.

In other preferred embodiments: the second nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000,
30 6000, or 8060 nucleotides in length; the second nucleic acid is a full length retroviral genome.

In another aspect, the invention features a method of screening a human subject for the presence or expression of an endogenous porcine retrovirus or retroviral sequence comprising:

35 contacting a target nucleic acid derived from the human subject with a second sequence chosen from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the

sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from
5 nucleotides 2452-4839 (e.g, from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g, from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or
10 antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of
15 SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid under conditions in which the sequences can hybridize, hybridization being indicative of the presence of the endogenous porcine retrovirus or retroviral sequence in the human subject.

20 In preferred embodiments, the target nucleic acid derived from a human subject is DNA, RNA or cDNA sample, nucleic acid from a blood sample or a tissue sample, e.g., a tissue biopsy sample.

In preferred embodiments, the human subject is a miniature swine or swine xenograft recipient, or a person who has come into contact with a miniature swine or swine
25 xenograft recipient.

In preferred embodiments, the second nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3
30 or its complement.

In other preferred embodiments: the second nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000,
35 6000, or 8060 nucleotides in length; the second nucleic acid is a full length retroviral genome.

In preferred embodiments: the recipient is tested for the presence of porcine retroviral sequences prior to implantation of swine or miniature swine tissue.

In another aspect, the invention features a method of screening for viral mutations which modulate, e.g., increase or decrease, susceptibility of a porcine retrovirus to an antiviral agent, e.g., an antiviral antibiotic. The method includes:

- administering a treatment, e.g., an antiviral agent, e.g., an antiviral antibiotic;
- 5 isolating a putative mutant porcine retroviral strain;
- determining a structure of the putative mutant retroviral strain; and
- comparing the structure to SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In another aspect, the invention features a method of screening for viral mutations which modulate, e.g., increase or decrease, susceptibility of a porcine retrovirus to an antiviral agent, e.g., an antiviral antibiotic. The method includes:

- growing the porcine retrovirus in a presence of a treatment, e.g., an antiviral agent, e.g., an antiviral antibiotic; and
- determine the amount of porcine retroviral DNA synthesized by hybridizing the
- 15 porcine retroviral DNA to a second sequence chosen from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a
- sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its
- 20 complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g, from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g, from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or
- 25 naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive
- 30 nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a
- 35 Tsukuba nucleic acid.

In preferred embodiments, the method further includes amplifying the porcine retroviral nucleic acid with primers which specifically hybridize to the sequence of SEQ ID

NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, e.g., by polymerase chain reaction quantitative DNA testing (PDQ).

In a preferred embodiment: the second nucleic acid is a Tsukuba-1 retroviral sequence, probe or primer, e.g., as described herein; the second nucleic acid is a porcine retroviral sequence, probe or primer, e.g., as described herein; the second nucleic acid is the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In preferred embodiments, the second nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In other preferred embodiments: the second nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the second nucleic acid is a full length retroviral genome.

In another aspect, the invention features a method for screening a porcine-derived product for the presence or expression of a swine or miniature swine retrovirus or retroviral sequence, e.g., an endogenous miniature swine retrovirus. The method includes:

contacting a target nucleic acid from the porcine-derived product with a second sequence chosen from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g., from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g., from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g., from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides

of sense or antisense sequence from nucleotides 2-1999 (e.g. from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g. from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid, under conditions in which hybridization can occur, hybridization being indicative of the presence or expression of an endogenous miniature swine or swine retrovirus or retroviral sequence s in the porcine-derived product.

In preferred embodiments the product is: a protein product, e.g., insulin; a food product; or a cellular transplant, e.g., a swine or miniature swine cell which is to be transplanted into a host, e.g., a swine or miniature swine cell which is genetically engineered to express a desired product,

In preferred embodiments, the method further includes amplifying the target nucleic acid with primers which specifically hybridize to the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In other preferred embodiments, the target nucleic acid is: DNA; RNA; or cDNA.

In preferred embodiments, the second nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In other preferred embodiments: the second nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the second nucleic acid is a full length retroviral genome.

In another aspect, the invention features a transgenic miniature swine or swine having a transgenic element, e.g., a base change, e.g., a change from A to G, or an insertion or a deletion of one or more nucleotides at an endogenous porcine retroviral insertion site, e.g., a retroviral insertion which corresponds to the retroviral genome of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In preferred embodiments, the transgenic element is a knockout, e.g., a deletion, insertion or a translocation, of one or more nucleic acids, which alters the activity of the endogenous porcine retrovirus.

In another aspect, the invention features a method of inhibiting expression of an endogenous porcine retrovirus, including: inserting a mutation, e.g. a deletion into the endogenous retrovirus.

In preferred embodiments, the endogenous porcine retrovirus is inactivated.

In preferred embodiments, the mutation can be a point mutation, an inversion, translocation or a deletion of one or more nucleotides of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

5 In another aspect, the invention features a method of detecting a recombinant virus or other pathogen, e.g., a protozoa or fungi. The method includes:

providing a pathogen having porcine retroviral sequence; and

determining if the pathogen includes non-porcine retroviral sequence, the presence of non-porcine retroviral sequence being indicative of viral recombination.

10 In preferred embodiments, the method further includes determining the structure of a retrovirus by comparing the retrovirus sequence with sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, a difference being indicative of viral recombination.

15 In preferred embodiments, the method further includes comparing the structure of the retrovirus with a human retroviral sequence, e.g., HTLV1, HIV1, or HIV2, a similarity in structure being indicative of viral recombination.

In another aspect, the invention features a method of determining the copy number, size, or completeness of a porcine retrovirus or retroviral sequence, e.g., in the genome of a donor, recipient or a graft. The method includes:

20 contacting a target nucleic acid from the donor, recipient or a graft, with a second sequence chosen from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10
25 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g. from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g. from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g. from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof;
30 a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof;
35 a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g. from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g. from nucleotides 4738-6722) of SEQ ID NO:2, or

nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid.

In preferred embodiments, the method further includes amplifying the porcine retroviral nucleic acid with primers which specifically hybridize to the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, e.g., by polymerase chain reaction quantitative DNA testing (PDQ) or nested PCR.

In preferred embodiments, the target nucleic acid includes: genomic DNA isolated from a miniature swine; RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine; DNA, RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine organ, e.g., a kidney; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a miniature swine organ which has been transplanted into a organ recipient, e.g., a xenogeneic recipient, e.g., a primate, e.g., a human.

In preferred embodiments, the target nucleic acid includes: genomic DNA isolated from a swine; RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine; DNA, RNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine organ, e.g., a kidney; RNA, DNA or cDNA, e.g., cDNA made from an RNA template, isolated from a swine organ which has been transplanted into a organ recipient, e.g., a xenogeneic recipient, e.g., a primate, e.g., a human.

In preferred embodiments, the second nucleic acid has at least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with a sequence from SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

In other preferred embodiments: the second nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length; the second nucleic acid is a full length retroviral genome.

In another aspect, the invention features a method for screening a tissue, e.g., a cellular or tissue transplant, e.g., a xenograft, or a tissue from a graft recipient, for the presence or expression of a swine or a miniature swine retroviral sequence, e.g., an endogenous miniature swine retrovirus. The method includes: contacting a tissue sample with an antibody specific for a retroviral protein, e.g., an anti-gag, pol, or env antibody, and thereby determining if the sequence is present or expressed.

In preferred embodiments the protein is encoded by a sequence from: the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

5 In preferred embodiments, the tissue is selected from the group consisting of: heart, lung, liver, bone marrow, kidney, brain cells, neural tissue, pancreas or pancreatic cells, thymus, or intestinal tissue.

A "purified preparation" or a "substantially pure preparation" of a polypeptide as used herein, means a polypeptide which is free from one or more other proteins, lipids, and nucleic acids with which it naturally occurs. Preferably, the polypeptide, is also separated
10 from substances which are used to purify it, e.g., antibodies or gel matrix, such as polyacrylamide. Preferably, the polypeptide constitutes at least 10, 20, 50 70, 80 or 95% dry weight of the purified preparation. Preferably, the preparation contains: sufficient polypeptide to allow protein sequencing; at least 1, 10, or 100 μ g of the polypeptide; at least 1, 10, or 100 mg of the polypeptide.

15 Specifically hybridize, as used herein, means that a nucleic acid hybridizes to a target sequence with substantially greater degree than it does to other sequences in a reaction mixture. By substantially greater means a difference sufficient to determine if the target sequence is present in the mixture.

A "treatment", as used herein, includes any therapeutic treatment, e.g., the
20 administration of a therapeutic agent or substance, e.g., a drug or irradiation.

A "purified preparation of nucleic acid", is a nucleic acid which is one or both of: not immediately contiguous with one or both of the coding sequences with which it is immediately contiguous (i.e., one at the 5' end and one at the 3' end) in the naturally-
25 occurring genome of the organism from which the nucleic acid is derived; or which is substantially free of a nucleic acid sequence or protein with which it occurs in the organism from which the nucleic acid is derived. The term includes, for example, a recombinant DNA which is incorporated into a vector, e.g., into an autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or eukaryote, or which exists as a separate molecule (e.g., a cDNA or a genomic DNA fragment produced by PCR or restriction
30 endonuclease treatment) independent of other DNA sequences. Substantially pure DNA also includes a recombinant DNA which is part of a hybrid gene encoding additional sequences. A purified retroviral genome is a nucleic acid which is substantially free of host nucleic acid or viral protein.

"Homologous", as used herein, refers to the sequence similarity between two
35 polypeptide molecules or between two nucleic acid molecules. When a position in both of the two compared sequences is occupied by the same amino acid or base monomer subunit, e.g., if a position in each of two DNA molecules is occupied by adenine, then the molecules are homologous at that position. The percent of homology between two sequences is a

function of the number of matching or homologous positions shared by the two sequences divided by the number of positions compared x 100. For example, if 6 of 10. of the positions in two sequences are matched or homologous then the two sequences are 60% homologous. By way of example, the DNA sequences ATTGCC and TATGGC share 50% homology. Generally, a comparison is made when two sequences are aligned to give maximum homology. The term sequence identity has substantially the same meaning.

The term "provirus" or "endogenous retrovirus," as used herein, refers to an integrated form of the retrovirus.

The terms "peptides", "proteins", and "polypeptides" are used interchangeably herein.

As used herein, the term "transgenic element" means a nucleic acid sequence, which is partly or entirely heterologous, i.e., foreign, to the animal or cell into which it is introduced but which is designed to be inserted, or is inserted, into the animal's genome in such a way as to alter the genome of the cell into which it is inserted. The term includes elements which cause a change in the sequence, or in the ability to be activated, of an endogenous retroviral sequence. Examples of transgenic elements include those which result in changes, e.g., substitutions (e.g., A for G), insertions or deletions of an endogenous retroviral sequence (or flanking regions) which result in inhibition of activation or misexpression of a retroviral product.

As used herein, the term "transgenic cell" refers to a cell containing a transgenic element.

As used herein, a "transgenic animal" is any animal in which one or more, and preferably essentially all, of the cells of the animal includes a transgenic element. The transgenic element can be introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection. This molecule may be integrated within a chromosome, or it may be extrachromosomally replicating DNA.

As described herein, one aspect of the invention features a pure (or recombinant) nucleic acid which includes a miniature swine (or swine) retroviral genome or fragment thereof, e.g., nucleotide sequence encoding a gag-pol or env polypeptide, and/or equivalents of such nucleic acids. The term "nucleic acid", as used herein, can include fragments and equivalents. The term "equivalent" refers to nucleotide sequences encoding functionally equivalent polypeptides or functionally equivalent polypeptides which, for example, retain the ability to react with an antibody specific for a gag-pol or env polypeptide. Equivalent nucleotide sequences will include sequences that differ by one or more nucleotide substitutions, additions or deletions, such as allelic variants, and will, therefore, include sequences that differ from the nucleotide sequence of gag, pol, or env shown in herein due to the degeneracy of the genetic code.

"Misexpression", as used herein, refers to a non-wild type pattern of gene expression, e.g., porcine retroviral, e.g., Tsukuba-1 gene expression, e.g., gag, pol or env gene expression. It includes: expression at non-wild type levels, i.e., over or under expression; a pattern of expression that differs from wild type in terms of the time or stage at which the gene is expressed, e.g., increased or decreased expression (as compared with wild type) at a predetermined developmental period or stage; a pattern of expression that differs from wild type in terms of decreased expression (as compared with wild type) in a predetermined cell type or tissue type; a pattern of expression that differs from wild type in terms of the splicing, size, amino acid sequence, post-translational modification, stability, or biological activity of the expressed, porcine retroviral, e.g., Tsukuba-1, polypeptides; a pattern of expression that differs from wild type in terms of the effect of an environmental stimulus or extracellular stimulus on expression of the porcine retroviral, e.g., Tsukuba-1 genes, e.g., a pattern of increased or decreased expression (as compared with wild type) in the presence of an increase or decrease in the strength of the stimulus.

Methods of the invention can be used with swine or miniature swine.

Endogenous retrovirus is a potential source of infection not always susceptible to conventional breeding practices. Many proviruses are defective and unable to replicate. Provirus, if intact, can be activated by certain stimuli and then initiate viral replication using the host's cellular mechanisms. Retroviral infection will often not harm the host cell. However, replication of virus may result in viremia, malignant transformation (e.g., via insertion of retroviral oncogenes), degeneration, or other insertional effects (e.g., gene inactivation). The effects of such infection may not emerge for many years. The spectrum of behavior of active lentiviral infection in humans is well described relative to HIV. These include AIDS, unusual infections and tumors, recombinant and other viruses, and antigenic variation which may prevent the generation of protective immunity by the infected host.

Screening of animals will allow elimination of donors with active replication of known viruses. Inactive proviruses can be detected with genetic probes and removed or inactivated. These novel approaches will allow the identification and elimination of potential human pathogens derived from swine in a manner not possible in the outbred human organ donor population and, thus, will be important to the development of human xenotransplantation.

The porcine retroviral sequences of the invention are also useful as diagnostic probes to detect activation of endogenous porcine retroviruses following transplantation and xenotransplantation of organs derived from swine or miniature swine. The porcine retroviral sequences of the invention also provide diagnostic tools necessary to assess the risks associated with transplantation of organs from swine or miniature swine into human recipients. These sequences are also useful for the longitudinal evaluation of retroviral activation in the human recipient of miniature swine-derived organs.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of cell biology, cell culture, molecular biology, transgenic biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are described in the literature. See, for example, *Molecular Cloning A Laboratory Manual*, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press: 1989); *DNA Cloning*, Volumes I and II (D. N. Glover ed., 1985); *Oligonucleotide Synthesis* (M. J. Gait ed., 1984); Mullis et al. U.S. Patent No. 4,683,195; *Nucleic Acid Hybridization* (B. D. Hames & S. J. Higgins eds. 1984); *Transcription And Translation* (B. D. Hames & S. J. Higgins eds. 1984); *Culture Of Animal Cells* (R. I. Freshney, Alan R. Liss, Inc., 1987); *Immobilized Cells And Enzymes* (IRL Press, 1986); B. Perbal, *A Practical Guide To Molecular Cloning* (1984); the treatise, *Methods In Enzymology* (Academic Press, Inc., N.Y.); *Gene Transfer Vectors For Mammalian Cells* (J. H. Miller and M. P. Calos eds., 1987, Cold Spring Harbor Laboratory); *Methods In Enzymology*, Vols. 154 and 155 (Wu et al. eds.), *Immunochemical Methods In Cell And Molecular Biology* (Mayer and Walker, eds., Academic Press, London, 1987); *Handbook Of Experimental Immunology*, Volumes I-IV (D. M. Weir and C. C. Blackwell, eds., 1986); *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986).

Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described below. All publications mentioned herein are incorporated by reference. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Detailed Description of the Drawings

Figure 1 is the nucleotide sequence (SEQ ID NO: 1) of the Tsukuba-1 cDNA.

Figure 2 is the nucleotide sequence (SEQ ID NO: 2) of a defective retroviral genome isolated from the retrovirus from the PK-15 cell line.

Figure 3 is the nucleotide sequence (SEQ ID NO: 3) of a retrovirus found in miniature swine.

Detailed Description

Miniature Swine Retroviruses

Transplantation may increase the likelihood of retroviral activation, if intact and infectious proviruses are present. Many phenomena associated with transplantation, e.g., immune suppression, graft rejection, graft-versus-host disease, viral co-infection, cytotoxic therapies, radiation therapy or drug treatment, can promote activation of retroviral expression.

Many species are thought to carry retroviral sequences in their genomic DNA. The number of intact (complete) retroviral elements that could be activated is often unknown.

Once activated, swine-derived viruses would require the appropriate receptor on human tissues to spread beyond the transplanted organ. Most intact endogenous proviruses (usually types B and C), once activated, are not pathogenic. However, coinfection with other viruses, recombination with other endogenous viruses, or modification of viral behavior in the foreign human environment may alter the pathogenicity, organ specificity or replication of the retroviruses or other infectious agents.

The lack of sequence data on pig viruses has impeded efforts to assess the number of porcine sequences, or porcine retroviral sequences, that have incorporated into the human genome or the frequency of incorporation.

The inventor, by showing that the Tsukuba-1 retrovirus is found in miniature swine, and by providing the entire sequence of the porcine retroviral (Tsukuba-1) genome, has allowed assessment of the risk of endogenous retroviruses in general clinical practice and more importantly in xenotransplantation.

The porcine retroviral sequences of the invention can be used to determine the level (e.g., copy number) of intact (i.e., potentially replicating) porcine provirus sequences in a strain of xenograft transplantation donors. For example, the copy number of the miniature swine retroviral sequences can be determined by the Polymerase Chain Reaction DNA Quantitation (PDQ) method, described herein, or by other methods known to those skilled in the art. This quantitation technique will allow for the selection of animal donors, e.g., miniature swine donors, without an intact porcine retroviral sequence or with a lower copy number of viral elements.

The porcine retroviral sequences of the invention can be used to determine if mutations, e.g., inversions, translocations, insertions or deletions, have occurred in the endogenous porcine retroviral sequence. Mutated viral genomes may be expression-deficient. For example, genetic lesions can be identified by exposing a probe/primer derived from porcine retrovirus sequence to nucleic acid of the tissue (e.g., genomic DNA) digested with a restriction endonucleases or by *in situ* hybridization of the probe/primer derived from the porcine retroviral sequence to the nucleic acid derived from donor, e.g., miniature swine, tissue. Alternatively, direct PCR analysis, using primers specific for porcine retroviral genes (e.g., genes comprising the nucleotide sequence shown in SEQ ID NO: 1, 2, or 3), can be used to detect the presence or absence of the genetic lesion in the porcine retroviral genome.

Miniature swine retroviral sequences of the invention can also be use to detect viral recombinants within the genome, or in the circulation, cells, or transplanted tissue, between the porcine retrovirus and other endogenous human viruses or opportunistic pathogens (e.g. cytomegalovirus) of the immunocompromised transplant recipient. For example, pieces of the viral genome can be detected via PCR or via hybridization, e.g., Southern or Northern

blot hybridization, using sequences derived from SEQ ID NO: 1, 2, or 3 as primers for amplification or probes for hybridization.

Miniature swine retroviral sequences of the invention, e.g., PCR primers, allow quantitation of activated virus. Sequences of the invention also allow histologic
5 localization (e.g., by *in situ* hybridization) of activated retrovirus. Localization allows clinicians to determine whether a graft should be removed as a source of potential retroviral infection of the human host or whether the retroviral infection was localized outside the graft. Sequences of the invention, e.g., PCR primers, allow the detection of actively
10 replicating virus, e.g., by using reverse transcribed PCR techniques known in the art. Standard techniques for reverse transcriptase measurements are often complicated, species-specific, and are of low sensitivity and specificity, and false positive results may develop using full-length probes for Southern and Northern molecular blotting. Sequences of the invention allow for sensitive and specific assays for the activation of virus and this will allow performance of a wide variety of tests, some of which are outlined below.

15 The invention provides for the testing and development of donor animals having reduced numbers of intact proviral insertions. It also provides for the testing of immunosuppressive regimens less likely to provide the conditions for active replication of retrovirus. Conditions likely to activate one retrovirus are generally more likely to activate other viruses including unknown retroviruses and known human pathogens including
20 cytomegalovirus, hepatitis B and C viruses, Human Immunodeficiency Viruses (I and II). Given the availability of preventative therapies for these infections, these therapies could be used prophylactically in patients known to be susceptible to the activation of porcine retrovirus.

The miniature swine retroviral sequences of the invention can be used to measure
25 the response of the miniature swine retroviral infection in humans to therapy, e.g., immunomodulatory or antiviral therapy, e.g., antiviral agents, e.g., antiviral antibiotics. With HIV, susceptibility to antiviral antibiotics is determined by the genetic sequence of the reverse transcriptase gene (RT pol region) and other genes. The ability to determine the exact sequence of the retroviral genes will allow the detection of mutations occurring
30 during infection which would then confer resistance of this virus to antiviral agents. Primers, e.g., for the RT-pol region, of the invention can be used to detect and to sequence clinical viral isolates from patients which have developed mutations by PDQ method described herein. The primers of the invention can also be used to determine whether
35 tumor cells, e.g., cancer cells, e.g. lymphoma or hepatocellular carcinoma, developing in xenograft recipients contain porcine retroviral elements.

The porcine retroviral sequences of the invention can also be used to detect other homologous retroviruses and to determine whether these are the same or different as compared to the Tusukuba-1 retroviral sequences. For example, within a species, the

polymerase genes are highly conserved. PCR assays aimed at the gag-pol region followed by sequence analysis allow for this detection of homologous viruses. The appropriate regions of the Tsukuba-1 virus can be determined by using sequences derived from SEQ ID NO:1, described herein, to identify additional 5' and 3' viral genomic sequences. As is discussed elsewhere herein, the sequences from SEQ ID NO: 1 were used to obtain the sequence of the PK-15 retroviral insert (SEQ ID NO:2) and of a retroviral insertion in a miniature swine (SEQ ID NO:3).

Miniature swine retroviral sequences of the invention can be used to screen donor animals and xenograft recipients after transplantation both for infection, and as a measure of the appropriate level of immune suppression, regarding susceptibility to infection. Physicians, medical staff, family, or individuals who come into contact with graft recipients, and others, can be screened for infection with virus derived from the xenograft recipient. Members of the population in general can also be screened. Such screening can be used for broad epidemiologic studies of the community. These methods can help in meeting the requirements of the F.D.A. regarding enhancing the safety of the recipients and of the community to exposure to new viruses introduced into the community by xenograft transplantation.

As is shown in Suzuka et al., 1986, FEBS 198:339, the swine retroviruses such as the Tsukuba-1 genome can exist as a circular molecule. Upon cloning the circular molecule is generally cleaved to yield a linear molecule. As will be understood by one skilled in the art, the start point and end point of the resulting linear molecule, and the relative subregions of the viral sequence will of course vary with the point of cleavage. For example, in the Suzuka et al. reference the LTR is shown to be in an internal fragment. This is indicated herein in that the order of gag, pol, env in SEQ ID NO 1 is shown as env, gag, pol, while elsewhere herein the order of these regions is given as the naturally occurring gag, pol, env order.

Primers Derived from the Porcine Retroviral (Tsukuba-1) Genome Sequence

A number of different primers useful in the methods of the invention have been described herein. One skilled in the art can identify additional primers from the viral sequence of SEQ ID NO:1 by using methods known in the art. For example, when trying to identify potentially useful primers one skilled in the art would look for sequences (sequences should be between about 15 and 30 nucleotides in length) which hybridize to SEQ ID NO:1 with high melting temperature; have a balanced distribution of nucleotides, e.g., a balanced distribution of A, T, C and Gs; have a terminal C or G; do not self-hybridize or internally complement.

Use of Primers Derived from the Porcine Retroviral (Tsukuba-1) Genome Sequence

I. Testing of organs or cells prior to transplantation

Potential donor animals can be screened for active retroviral replication prior to being used in transplantation. This allows avoidance of animals undergoing active viral replication. Replicating virus is often infectious in 100% of recipients, while nonreplicating, latent provirus generally causes infection in 5 to 25% of recipients.

5 II. Testing of recipients

Serial samples, e.g., of white blood cells, can be obtained from a graft recipient monthly, e.g., for the first month and every three months thereafter. Tissue biopsies obtained for evaluation of graft function can be used to evaluate the activation of retroviral sequences or of the expression of retroviral sequences in graft tissue. Samples can be screened
10 for the presence of retrovirus infection both specifically for the homologous virus, for viral recombinants containing portions of the viral genome, and for other retroviruses, using, e.g., PCR primers for the pol region of the virus, which is the region most likely to be conserved. If virus is detected, quantitative PCR can be used to determine the relative stability of viral production. Cells isolated from xenograft recipients can be tested by
15 cocultivation with permissive human and porcine (e.g., pig fallopian tube, pig macrophage, or pig testis) cell lines known to contain endogenous viruses. Isolated virus will be tested for homology with the parental strain and for mutations which might affect susceptibility to antiviral agents, e.g., antiviral antibiotics.

20 III. Testing of surgical and medical personnel and family members of graft recipient

Samples, e.g., white blood cells, can be banked (archived) from the surgical and medical personnel and from family members of the recipient prior to transplantation and at three months intervals for the first year and at least annually thereafter. Epidemiologic studies can be performed on these samples as well. These samples can be tested if the
25 recipient becomes viremic or if unusual clinical manifestations are noted in these individuals.

IV. Testing of tumor cells

Tumor cells which develop from a graft, or a graft recipient, can be tested for the presence of active retrovirus and for proviruses.

30 V. Testing of patients

Patients can be retested for any significant change in clinical condition or for increased immune suppression of graft rejection which may be associated with an increased risk of viral activation.

Sequencing of the porcine retroviral (Tsukuba-1) genome

35 A clone (P λ 8.8) containing the 8060 bp XhoI porcine retrovirus (Tsukuba-1) insert was used to transfect competent *E. coli*, and DNA was isolated for sequencing. The strategy used to sequence the 8060 bp porcine retrovirus genome included a combination of procedures which are outlined below.

Random fragments (1-3 kb) of the clone (Pλ8.8) were generated by sonication. The fragments were blunt-ended and were subcloned into the EcoRV site of the pBluescript SK vector. Plasmid DNA was prepared using a modified alkaline lysis procedure. DNA sequencing was performed using DyeDeoxy termination reactions (ABI). Base specific fluorescent dyes were used as labels. Sequencing reactions were analyzed on 4.75% polyacrylamide gels by an ABI 373A-S or 373S automated sequencer. Subsequent data analysis was performed on Sequencer™ 3.0 software. The following internal sequencing primers were synthesized:

| | | | | | |
|----|------|----|------------------------|----|----------------|
| 10 | AP1 | 5' | GATGAACAGGCAGACATCTG | 3' | (SEQ ID NO:48) |
| | AP2 | 5' | CGCTTACAGACAAGCTGTGA | 3' | (SEQ ID NO:49) |
| | AP3 | 5' | AGAACAAAGGCTGGGAAAGC | 3' | (SEQ ID NO:50) |
| | AP4 | 5' | ATAGGAGACAGCCTGAACTC | 3' | (SEQ ID NO:51) |
| | AP5 | 5' | GGACCATGTCTGACCCTAT | 3' | (SEQ ID NO:52) |
| 15 | AP6 | 5' | GTCAACACCTATAACCAGCTC | 3' | (SEQ ID NO:53) |
| | AP7 | 5' | CATCTGAGGTATAGCAGGTC | 3' | (SEQ ID NO:54) |
| | AP8 | 5' | GCAGGTGTAGGAACAGGAAC | 3' | (SEQ ID NO:55) |
| | AP9 | 5' | ACCTGTTGAACCATCCCTCA | 3' | (SEQ ID NO:56) |
| | AP10 | 5' | CGAATGGAGAGATCCAGGTA | 3' | (SEQ ID NO:57) |
| 20 | AP11 | 5' | CCTGCATCACTTCTCTTACC | 3' | (SEQ ID NO:58) |
| | AP12 | 5' | TTGCCTGCTTGTGGAATACG | 3' | (SEQ ID NO:59) |
| | AP13 | 5' | CAAGAGAAGAAGTGGGGAATG | 3' | (SEQ ID NO:60) |
| | AP14 | 5' | CACAGTCGTACACCACGCAG | 3' | (SEQ ID NO:61) |
| | AP15 | 5' | GGGAGACAGAAGAAGAAAGG3' | | (SEQ ID NO:62) |
| 25 | AP16 | 5' | CGATAGTCATTAGTCCCAGG | 3' | (SEQ ID NO:63) |
| | AP17 | 5' | TGCTGGTTTGCATCAAGACCG | 3' | (SEQ ID NO:64) |
| | AP18 | 5' | GTCGCAAAGGCATACCTGCT | 3' | (SEQ ID NO:65) |
| | AP19 | 5' | ACAGAGCCTCTGCTAAGAAG | 3' | (SEQ ID NO:66) |
| | AP20 | 5' | GCAGCTGTTGACAATCATC | 3' | (SEQ ID NO:67) |
| 30 | AP21 | 5' | TATGAGGAGAGGGCTTGACT | 3' | (SEQ ID NO:68) |
| | AP22 | 5' | AGCAGACGTGCTAGGAGGT | 3' | (SEQ ID NO:69) |
| | AP23 | 5' | TCCTCTTGCTGTTTGCATC | 3' | (SEQ ID NO:70) |
| | AP24 | 5' | CAGACACTCAGAACAGAGAC | 3' | (SEQ ID NO:71) |
| | AP25 | 5' | ACATCGTCTAACCCACCTAG | 3' | (SEQ ID NO:72) |
| 35 | AP26 | 5' | CTCGTTTCTGGTCATACCTGA | 3' | (SEQ ID NO:73) |
| | AP27 | 5' | GAGTACATCTCTCTAGGCA | 3' | (SEQ ID NO:74) |
| | AP28 | 5' | TGCCTAGAGACATGTACTC | 3' | (SEQ ID NO:4) |
| | AP29 | 5' | CCTCTTCTAGCCATTCTTCA | 3' | (SEQ ID NO:5) |

40 The clone (Pλ8.8) containing the 8060 bp XhoI porcine retrovirus (Tsukuba-1) insert was deposited with ATCC on December 27, 1995 (ATCC Deposit No.97396).

Determination of the porcine retroviral (Tsukuba-1) copy number in a miniature swine

Total genomic DNA was isolated from miniature swine kidney by the methods known in the art. The isolated genomic DNA was digested with either EcoRI or HindIII restriction enzyme. The DNA digests were electrophoresed on an agarose gel, Southern blotted and hybridized to the full-length, purified, Tsukuba-1 sequence (SEQ ID NO:1) under high stringency conditions (0.1 X SSC, 65°C). In both digested samples (EcoRI or HindIII) at least six copies of the high molecular fragments of the miniature swine genome

(over 16 Kb in size) hybridized to SEQ ID NO:1, indicating the presence of homologous retroviral sequences in porcine DNA.

Susceptibility Testing by Polymerase Chain Reaction DNA Quantitation (PDQ)

5 Polymerase chain reaction (PCR) DNA quantitation (PDQ) susceptibility testing can be used to rapidly and directly measure nucleoside sensitivity of porcine retrovirus isolates.

PCR can be used to quantitate the amount of porcine retroviral RNA synthesized after *in vitro* infection of peripheral blood mononuclear cells. The relative amounts of porcine retroviral RNA in cell lysates from cultures maintained at different drug concentrations reflect drug inhibition of virus replication. With the PDQ method both infectivity titration and susceptibility testing can be performed on supernatants from primary cultures of peripheral blood mononuclear cells.

10 The PDQ experiments can be performed essentially as described by Eron et al., *PNAS USA* 89:3241-3245, 1992. Briefly, aliquots (150 μ l) of serial dilutions of virus sample can be used to infect 2×10^6 PHA-stimulated donor PBMCs in 1.5 ml of growth medium per well of a flat-bottom 24-well plate (Corning). Separate cell samples can be counted, harvested, and lysed at 48, 72 and 96 hr. Quantitative PCR and porcine retrovirus copy-number determination can then be performed in duplicate on each lysate.

20 The results of a PDQ infectivity titration assay can be used to determine the virus dilution and length of culture time employed in a subsequent PDQ susceptibility test. These parameters should be chosen so that the yield of porcine retrovirus specific PCR product for the untreated control infection would fall on the porcine retrovirus copy-number standard curve before the curve approached its asymptotic maximum, or plateau. PHA-stimulated donor PBMCs can be incubated with drug for 4 hr prior to infection. Duplicate wells in a 24-well plate should receive identical porcine retrovirus inocula for each drug concentration tested and for the untreated infected controls. Uninfected controls and drug toxicity controls should be included in each experiment. All cultures can be harvested and cells lysed for PCT after either 48 or 72 hr. Previously characterized isolates can be used as assay standards in each experiment.

30 Cell pellets can be lysed in various volumes of lysis buffer (50 mM KCl/10mM Tris •HCl, pH 8.3/2.5 mM MgCl₂/0.5% Nonidet P-40/0.5% Tween 20/0.01% proteinase K) to yield a concentration of 1.2×10^4 cell equivalents/ μ l. Uniformity to cell lysate DNA concentrations should be confirmed in representative experiments by enhancement of Hoechst 33258 fluorescence (Mini-Fluorometer, Hoefer).

35 A conserved primer pair can be synthesized according to the pol gene sequences. The primers can then be used to amplify a 1580-base pair fragment of the porcine retrovirus pol gene from 1.2×10^5 cell equivalents of lysate by using PCR (GeneAmp, Cetus) under

standard conditions. Amplifications should be repeated if porcine retrovirus DNA is amplifiable from reagent controls.

Porcine retrovirus pol gene amplification products can be specifically detected and quantitated as described (Conway, B.C. (1990) in Techniques in HIV Research, (Aldovani & Walker, eds.) (Stockton, New York) pp.40-46). Heat-denatured PCR products can be hybridized in a Streptavidin-coated microtiter plate well with both biotinylated capture probe and horseradish peroxidase (HRP)-labeled detector probe [enzyme-linked oligonucleotide solution sandwich hybridization assay ((ELOSAs), DuPont Medical Products, Billerica, MA) for 60 min at 37°C. After extensive washing to remove all reactants except probe-DNA hybrids, an HRP chromogen, tetramethylbenzidine (TMBBlue, Transgenic Sciences, Worcester, MA), should be added to each well. The HRP-catalyzed color development should be stopped after 1 hr by addition of sulfuric acid to 0.65 M. Absorbance (OD) at 450 nm can be measured in an automated microtiter plate reader (SLT Lab Instruments, Hillsborough, NC).

A standard curve of porcine retrovirus DNA copy number can be generated in each PCR by using a dilution series of cells containing one porcine proviral genome per cell. Preparation of a miniature swine having a knockout of Tsukuba-1 viral sequence using isogenic DNA targeting vectors

Isogenic DNA, or DNA that is substantially identical in sequence between the targeting vector and the target DNA in the chromosomes, greatly increases the frequency for homologous recombination events and gene targeting efficiency. Using isogenic-DNA targeting vectors, targeting frequencies of 80% or higher can be achieved in mouse embryonic stem cells. This is in contrast to non-isogenic DNA vectors which normally yield targeting frequencies of around 0.5% to 5%, i.e., approximately two orders of magnitude lower than isogenic DNA vectors. Isogenic DNA constructs are predominantly integrated into chromosomes by homologous recombination rather than random integration. As a consequence, targeted mutagenesis of viral sequences, e.g., viral genes, can be carried out in biological systems including zygotes, which do not lend themselves to the use of elaborate selection protocols, resulting in production of animals, e.g., miniature swine, free of, or having a reduced number of, activatable viral sequences. In order for the isogenic DNA approach to be feasible, targeting vectors should be constructed from a source of DNA that is identical to the DNA of the organism to be targeted. Ideally, isogenic DNA targeting is carried out in inbred strains of animals, e.g., inbred miniature swine, in which all genetic loci are homozygous. Any animal of that strain can serve as a source for generating isogenic targeting vectors. This protocol for isogenic gene targeting is outlined in TeRiele et al., PNAS 89:5128-5132, 1992 and PCT/US92/07184, herein incorporated by reference. A protocol for producing Tsukuba-1 knockout miniature swine is described briefly below.

An insertion vector is designed as described by Hasty and Bradley (Gene Targeting Vectors for Mammalian Cells, in Gene Targeting: A Practical Approach, ed. Alexandra L. Joyner, IRL Press 1993). Insertion vectors require that only one crossover event occur for integration by homologous recombination into the native locus. The double strand breaks, the two ends of the vector which are known to be highly recombinogenic, are located on adjacent sequences on the chromosome. The targeting frequencies of such constructions will be in the range of 30 to 50%. One disadvantage of insertion vectors, in general, concerns the sequence duplications that are introduced and that potentially make the locus unstable. All these constructions are made using standard cloning procedures.

Replacement vectors have also been extensively described by Hasty and Bradley. Conceptually more straight forward than the insertion vector, replacement vectors use an essentially co-linear fragment of a stretch of Tsukuba-1 genomic sequence. Preferably, the DNA sequence from which an isogenic replacement vector is constructed includes approximately 6 to 10 kb of uninterrupted DNA. Two crossovers, one on either side of the selectable marker causes the mutant targeting vector to become integrated and replace the wild-type gene.

Microinjection of the isogenic transgene DNA into one of the pronuclei of a porcine embryo at the zygote stage (one-cell embryo) is accomplished by modification of a protocol described earlier (Hammer et al. 1985, Nature 315, 680; Pursel et al. 1989, Science 244, 1281). The age and the weight of the donor pigs, e.g., haplotype specific mini-swine, are critical to success. Optimally, the animals are of age 8 to 10 months and weigh 70 to 85 lbs. This increases the probability of obtaining an adequate supply of one-cell embryos for microinjection of the transgenes. In order to allow for accurate timing of the embryo collections at this stage from a number of embryo donors, the gilts are synchronized using a preparation of synthetic progesterone (Regumate). Hormone implants are applied to designated gilts 30 days prior to the date of embryo collection. Twenty days later, ten days prior to the date of collection, the implants are removed and the animals are treated with additional hormones to induce superovulation to increase the number of embryos for microinjection. Three days following implant removal, the animals are treated with 400 to 1000 IU of pregnant mare serum gonadotropin (PMSG) and with 750 IU of human chorionic gonadotropin (hCG) three to four days later. These animals are bred by artificial insemination (AI) on two consecutive days following injection of hCG.

Embryo collections are performed as follows: three days following the initial injection of hCG, the animals are anesthetized with an intramuscular injection of Telazol (3 mg/lb), Rompum (2 mg/lb) and Atropine (1 mg/lb). A midline laparotomy is performed and the reproductive tract exteriorized. Collection of the zygotes is performed by cannulating the ampulla of the oviduct and flushing the oviduct with 10 to 15 ml phosphate buffered saline, prewarmed to 39° C. Following the collection the donor animals are

prepared for recovery from surgery according to USDA guidelines. Animals used twice for embryo collections are euthanized according to USDA guidelines.

Injection of the transgene DNA into the pronuclei of the zygotes is carried out as summarized below: Zygotes are maintained in medium HAM F-12 supplemented with 10% fetal calf serum at 38° C in 5% CO₂ atmosphere. For injection the zygotes are placed into BMOC-2 medium, centrifuged at 13,000 g to partition the embryonic lipids and visualize the pronuclei. The embryos are placed in an injection chamber (depression slide) containing the same medium overlaid with light paraffin oil. Microinjection is performed on a Nikon Diaphot inverted-microscope equipped with Nomarski optics and Narishige micromanipulators. Using 40x lens power the embryos are held in place with a holding pipette and injected with a glass needle which is back-filled with the solution of DNA containing the transgenic element, e.g., a mutant viral gene (2 µg/ml). Injection of approximately 2 picoliters of the solution (4 femtograms of DNA), which is equivalent to around 500 copies of the transgenic element, e.g., a mutant viral gene, is monitored by the swelling of the pronucleus by about 50%. Embryos that are injected are placed into the incubator prior to transfer to recipient animals.

Recipient animals are prepared similarly to the donor animals, but not superovulated. Prior to the transfer of the injected embryos, recipient gilts are anesthetized, the abdomen opened surgically by applying a longitudinal incision and the ovaries exteriorized. The oviduct ipsilateral to the ovary with the larger number of corpus lutei is flushed, the embryos checked to evaluate if the animals is reproductively sound. Approximately 4 to 6 zygotes injected with the transgenic element, e.g., a mutant viral gene, are transferred to the flushed oviduct, the abdominal incision sutured and the animals placed in a warm area for recovery. The status of the pregnancy is monitored by ultrasound starting at day 25, or approximately one week following the expected date of implantation. Pregnant recipients are housed separately until they are due to farrow.

Newborn piglets are analyzed for integration of the transgenic element into chromosomal DNA. Genomic DNA is extracted from an ear punch or a blood sample and initial screening is performed using PCR. Animals that are potentially transgenic element-positive are confirmed by Southern analysis. Transgenic founder animals are subjected to further analysis regarding the locus of transgenic element integration using Southern analysis.

The isolation and sequencing of an endogenous swine retroviral insert and of a retroviral insert in porcine PK-15 cells

Cloning of PK15 and PAL endogenous retroviruses

I. Poly A⁺ RNA isolation

Peripheral blood lymphocytes (PBLs) were prepared from haplotype d/d miniswine using standard protocols known in the art. The PBLs were cultured in the presence of 1% phytohemagglutinin (PHA) for about 84 hours. The activated PBLs were collected and total RNA was isolated using commercially available kits, such as Gentra's (Minneapolis, Minnesota) PUREscript Kit. Poly A+RNA was isolated from the total RNA using another commercially available product, Dynal Dynabeads (Lake Success, NY). Northern analysis of the RNA using a pig retroviral probe confirmed the presence of potentially full-length retroviral genome RNA. RNA from PK15 cells was isolated using similar protocols.

10 II. Construction of the cDNA libraries

Using Superscript Choice System (Life Technologies Ltd, Gibco BRL, Gaithersburg, MD) for cDNA Synthesis, a cDNA library was constructed using oligo dT to make the first strand cDNA. The use of Superscript reverse transcriptase was important in order to obtain full-length retroviral (RV) cDNAs, due to the length of the RV RNA. The cDNA library was enriched for large cDNA fragments by size selecting >4 kb fragments by gel electrophoresis. The cDNAs were cloned into Lambda ZAP Express (Clontech Laboratories, Inc. Palo Alto, CA), which is one of the few commercially available cDNA vectors that would accept inserts in the 1-12kb range.

20 III. Screening of the cDNA libraries

0.75 - 1.2 x 10⁶ independent clones were screened using either gag and pol or gag and env probes. Double positive clones were further purified until single isolates were obtained (1 or 2 additional rounds of screening).

25 IV. Characterization of the clones

Between 18 and 30 double positive clones were selected for evaluation. Lambda DNA was prepared using standard protocols, such as the Lambda DNA Kit (Qiagen Inc., Chatsworth, CA). The clones were analyzed by PCR to check for (a) RV genes, and (b) determine the size of insert and LTR regions. Restriction digests were also done to confirm the size of insert and to attempt to categorize the clones. Clones containing the longest inserts and having consistent and predicted PCR data were sequenced.

Development of a PCR-based assay for the detection of the presence of an endogenous retrovirus in cells, tissues, organs, miniswine or recipient hosts (e.g., primates, humans)

Using a commercially available computer software program (such as RightPrimer, Oligo 4.0, MacVector or Geneworks), one can analyze sequences disclosed herein for the selection of PCR primer pairs. The criteria for the general selection of primer pairs includes:

- a. The T_m of each primer is between 65-70°C

- b. The T_m 's for each pair differ by no more than 3°C
- c. The PCR fragment is between 200-800 bp in length
- d. There are no repeats, self complementary bases, primer-dimer issues, etc for each pair

5

A. Additional criteria for: A pig-specific PCR assay

- a. Primers are selected within porcine-specific regions of the sequence -- such as within gag, env, or U3. Porcine-specific primers are defined as sequences which overall have <70% homology to the corresponding region in human, mouse and primate retroviruses. In addition, the last five bases at the 3' end of the primer should be unique to the pig retroviral sequence.

10

- b. Primers should have no more than one or two mismatched bases based on the miniswine, and retroviral sequences disclosed herein. These mismatched bases should not be within the last three or four bases of the 3' end of the primer.

15

B. Additional criteria for: Miniswine-specific PCR assay

- a. Primers are selected such that there are at least one or two mismatches between miniswine and domestic pig sequences. At least one of these mismatches should be located within the last three or four bases at the 3' end of the primer. Preferably, these mismatches would be a change from either a G or C in miniswine to either an A or T in domestic pig.

20

RT-PCR Strategy

There are a number of commercially available RT-PCR Kits for routine amplification of fragments. Several primer pairs should be tested to confirm T_m and specificity. Location of primers within the sequence depends in part on what question is being answered. RT-PCR should answer questions about expression and presence of RV sequences. PCR will not necessarily answer the question of whether the retroviral sequence is full-length or encodes a replication competent retrovirus. A positive signal in these tests only says there is RV sequence present. Indication of the possibility of full-length viral genomes being present can be obtained by performing long PCR using primers in U5 and U3. A commercial kit for long RT-PCR amplification is available (Takara RNA LA PCR Kit). Confirmation of full-length viral genomes requires infectivity studies and/or isolation of viral particles.

25

30

Northern analyses would complement RT-PCR data. Detection of bands at the predicted size of full-length viral genomes with hybridization probes from env, U3 or U5 would provide stronger evidence. The presence of other small bands hybridizing would indicate the amount of defective viral fragments present.

35

Elisa-Based Assay To Detect The Presence Of Porcine Retroviral Proteins, Polypeptides Or Peptides

In addition to the use of nucleic acid-based, e.g., PCR-based assays, to detect the presence of retroviral sequences, ELISA based assays can detect the presence of porcine retroviral proteins, polypeptides and peptides.

The basic steps to developing an ELISA include (a) generation of porcine retroviral specific peptides, polypeptides and proteins; (b) generation of antibodies which are specific for the porcine retroviral sequences; (c) developing the assay.

Using the retroviral sequences disclosed herein, antigenic peptides can be designed using computer based programs such as MacVector or Geneworks to analyse the retroviral sequences. Alternatively, it is possible to express the porcine retroviral sequences in gene expression systems and to purify the expressed polypeptides or proteins. After synthesis, the peptides, polypeptides or proteins are used to immunize mice or rabbits and to develop serum containing antibodies.

Having obtained the porcine retroviral specific antibodies the ELISA can be developed as follows. ELISA plates are coated with a volume of polyclonal or monoclonal antibody (capture antibody) which is reactive with the analyte to be tested. Such analytes include porcine retroviruses or retroviral proteins such as env or p24. The ELISA plates are then incubated at 4°C overnight. The coated plates are then washed and blocked with a volume of a blocking reagent to reduce or prevent non-specific hybridization. Such blocking reagents include bovine serum albumin (BSA), fetal bovine serum (FBS), milk, or gelatin. The temperature for the blocking process is 37°C. Plates can be used immediately or stored frozen at -20°C until needed. The plates are then washed, loaded with a serial dilution of the analyte, incubated at 37°C, and washed again. Bound analyte is detected using a detecting antibody. Detecting antibodies include enzyme-linked, fluoresceinated, biotin-conjugated or other tagged polyclonal or monoclonal antibodies which are reactive with the analyte. If monoclonal antibodies are used the detecting antibody should recognize an epitope which is different from the capture antibody.

Other Embodiments

In another aspect, the invention provides a substantially pure nucleic acid having, or comprising, a nucleotide sequence which encodes a swine or miniature swine, e.g., a Tsukuba-1 retroviral gag polypeptide.

In preferred embodiments: the nucleic acid is or includes the nucleotide sequence from nucleotides 2452-4839 of SEQ ID NO:1; the nucleic acid is at least 60%, 70%, 80%, 90%, 95%, 98%, or 99% homologous with a nucleic acid sequence corresponding to nucleotides 2452-4839 of SEQ ID NO:1; or by a sequence which, hybridizes under high stringency conditions to nucleotides 2452-4839 of SEQ ID NO:1; the nucleic acid includes a fragment of SEQ ID NO:1 which is at least 25, 50, 100, 200, 300, 400, 500, or 1,000

bases in length; the nucleic acid differs from the nucleotide sequence corresponding to nucleotides 2452-4839 of SEQ ID NO:1 due to degeneracy in the genetic code; the nucleic acid differs from the nucleic acid sequence corresponding to nucleotides 2452-4839 of SEQ ID NO:1 by at least one nucleotide but by less than 5, 10, 15 or 20 nucleotides and
5 preferably which encodes an active peptide.

In yet another preferred embodiment, the nucleic acid of the invention hybridizes under stringent conditions to a nucleic acid probe corresponding to at least 12 consecutive nucleotides from nucleotides 2452-4839 of SEQ ID NO:1, or more preferably to at least 20 consecutive nucleotides from nucleotides 2452-4839 of SEQ ID NO:1, or more preferably
10 to at least 40 consecutive nucleotides from nucleotides 2452-4839 of SEQ ID NO:1.

In another aspect, the invention features, a purified recombinant nucleic acid having at least 50%, 60%, 70%, 80%, 90%, 95%, 98%, or 99% homology with a nucleotide sequence corresponding to nucleotides 2452-4839 of SEQ ID NO:1.

The invention also provides a probe or primer which includes or comprises a
15 substantially purified oligonucleotide. The oligonucleotide includes a region of nucleotide sequence which hybridizes under stringent conditions to at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 of SEQ ID NO:1, or naturally occurring mutants thereof. In preferred embodiments, the probe or primer further includes a label attached thereto. The label can be, e.g., a radioisotope, a fluorescent compound, an
20 enzyme, and/or an enzyme co-factor. Preferably the oligonucleotide is at least 10 and less than 20, 30, 50, 100, or 150 nucleotides in length. Preferred primers of the invention include oligonucleotides having a nucleotide sequence shown in any of SEQ ID NOs:32-37.

The invention involves nucleic acids, e.g., RNA or DNA, encoding a polypeptide of
25 the invention. This includes double stranded nucleic acids as well as coding and antisense single strands.

In another aspect, the invention provides a substantially pure nucleic acid having, or comprising, a nucleotide sequence which encodes a swine or miniature swine, e.g., a Tsukuba-1 retroviral pol polypeptide.

In preferred embodiments: the nucleic acid is or includes the nucleotide sequence
30 corresponding to nucleotides 4871-8060 of SEQ ID NO:1; the nucleic acid is at least 60%, 70%, 80%, 90%, 95%, 98%, or 99% homologous with a nucleic acid sequence corresponding to nucleotides 4871-8060 of SEQ ID NO:1; or by a sequence which, hybridizes under high stringency conditions to nucleotides 4871-8060 of SEQ ID NO:1; the
35 nucleic acid includes a fragment of SEQ ID NO:1 which is at least 25, 50, 100, 200, 300, 400, 500, or 1,000 bases in length; the nucleic acid differs from the nucleotide sequence corresponding to nucleotides 4871-8060 of SEQ ID NO:1 due to degeneracy in the genetic code; the nucleic acid differs from the nucleic acid sequence corresponding to nucleotides

4871-8060 of SEQ ID NO:1 by at least one nucleotide but by less than 5, 10, 15 or 20 nucleotides and preferably which encodes an active peptide.

In yet another preferred embodiment, the nucleic acid of the invention hybridizes under stringent conditions to a nucleic acid probe corresponding to at least 12 consecutive
5 nucleotides from nucleotides 4871-8060 of SEQ ID NO:1, or more preferably to at least 20 consecutive nucleotides from nucleotides 4871-8060 of SEQ ID NO:1, or more preferably to at least 40 consecutive nucleotides from nucleotides 4871-8060 of SEQ ID NO:1.

In another aspect, the invention features, a purified recombinant nucleic acid having at least 50%, 60%, 70%, 80%, 90%, 95%, 98%, or 99% homology with a nucleotide
10 sequence corresponding to nucleotides 4871-8060 of SEQ ID NO:1.

The invention also provides a probe or primer which includes or comprises a substantially purified oligonucleotide. The oligonucleotide includes a region of nucleotide sequence which hybridizes under stringent conditions to at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, or naturally
15 occurring mutants thereof. In preferred embodiments, the probe or primer further includes a label attached thereto. The label can be, e.g., a radioisotope, a fluorescent compound, an enzyme, and/or an enzyme co-factor. Preferably the oligonucleotide is at least 10 and less than 20, 30, 50, 100, or 150 nucleotides in length. Preferred primers of the invention include oligonucleotides having a nucleotide sequence shown in any of SEQ ID NOs:38-
20 47.

The invention involves nucleic acids, e.g., RNA or DNA, encoding a polypeptide of the invention. This includes double stranded nucleic acids as well as coding and antisense single strands.

In another aspect, the invention provides a substantially pure nucleic acid having, or
25 comprising, a nucleotide sequence which encodes a swine or miniature swine, e.g., a Tsukuba-1 retroviral env polypeptide.

In preferred embodiments: the nucleic acid is or includes the nucleotide sequence corresponding to nucleotides 2-1999 of SEQ ID NO:1; the nucleic acid is at least 60%, 70%, 80%, 90%, 95%, 98%, or 99% homologous with a nucleic acid sequence
30 corresponding to nucleotides 2-1999 of SEQ ID NO:1; or by a sequence which, hybridizes under high stringency conditions to nucleotides 2-1999 of SEQ ID NO:1; the nucleic acid includes a fragment of SEQ ID NO:1 which is at least 25, 50, 100, 200, 300, 400, 500, or 1,000 bases in length; the nucleic acid differs from the nucleotide sequence corresponding to nucleotides 2-1999 of SEQ ID NO:1 due to degeneracy in the genetic code; the nucleic
35 acid differs from the nucleic acid sequence corresponding to nucleotides 2-1999 of SEQ ID NO:1 by at least one nucleotide but by less than 5, 10, 15 or 20 nucleotides and preferably which encodes an active peptide.

In yet another preferred embodiment, the nucleic acid of the invention hybridizes under stringent conditions to a nucleic acid probe corresponding to at least 12 consecutive nucleotides from nucleotides 2-1999 of SEQ ID NO:1, or more preferably to at least 20 consecutive nucleotides from nucleotides 2-1999 of SEQ ID NO:1, or more preferably to at least 40 consecutive nucleotides from nucleotides 2-1999 of SEQ ID NO:1.

In another aspect, the invention features, a purified recombinant nucleic acid having at least 50%, 60%, 70%, 80%, 90%, 95%, 98%, or 99% homology with a nucleotide sequence corresponding to nucleotides 2-1999 of SEQ ID NO:1.

The invention also provides a probe or primer which includes or comprises a substantially purified oligonucleotide. The oligonucleotide includes a region of nucleotide sequence which hybridizes under stringent conditions to at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 of SEQ ID NO:1, or naturally occurring mutants thereof. In preferred embodiments, the probe or primer further includes a label attached thereto. The label can be, e.g., a radioisotope, a fluorescent compound, an enzyme, and/or an enzyme co-factor. Preferably the oligonucleotide is at least 10 and less than 20, 30, 50, 100, or 150 nucleotides in length. Preferred primers of the invention include oligonucleotides having a nucleotide sequence shown in any of SEQ ID NOs:6-31.

The invention includes nucleic acids, e.g., RNA or DNA, encoding a polypeptide of the invention. This includes double stranded nucleic acids as well as coding and antisense single strands.

Included in the invention are: allelic variations, natural mutants, induced mutants, that hybridize under high or low stringency conditions to the nucleic acid of SEQ ID NO:1, 2, or 3 (for definitions of high and low stringency see Current Protocols in Molecular Biology, John Wiley & Sons, New York, 1989, 6.3.1 - 6.3.6, hereby incorporated by reference).

The invention also includes purified preparations of swine or miniature swine retroviral polypeptides, e.g., gag pol, or env polypeptides, or fragments thereof, preferably biologically active fragments, or analogs, of such polypeptides. In preferred embodiments: the polypeptides are miniature swine retroviruses polypeptides; the polypeptides are Tsukuba polypeptides; the polypeptides are gag, pol, or env polypeptides encoded by SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, or naturally occurring variants thereof.

A biologically active fragment or analog is one having any in vivo or in vitro activity which is characteristic of the Tsukuba-1 polypeptides described herein, or of other naturally occurring Tsukuba-1 polypeptides. Fragments include those expressed in native or endogenous cells, e.g., as a result of post-translational processing, e.g., as the result of the removal of an amino-terminal signal sequence, as well as those made in expression systems, e.g., in CHO cells. A useful polypeptide fragment or polypeptide analog is one

which exhibits a biological activity in any biological assay for Tsukuba-1 polypeptide activity. Most preferably the fragment or analog possesses 10%, preferably 40%, or at least 90% of the activity of Tsukuba-1 polypeptides, in any in vivo or in vitro Tsukuba-1 polypeptide assay.

5 In order to obtain a such polypeptides, polypeptide-encoding DNA can be introduced into an expression vector, the vector introduced into a cell suitable for expression of the desired protein, and the peptide recovered and purified, by prior art methods. Antibodies to the polypeptides can be made by immunizing an animal, e.g., a rabbit or mouse, and recovering antibodies by prior art methods.

10 The invention also features a purified nucleic acid, which has least 60%, 70%, 72%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, most preferably at least 98%, 99% or 100% sequence identity or homology with SEQ ID NO:1 or its complement, SEQ ID NO: 2 or its complement, or SEQ ID NO: 3 or its complement.

15 In preferred embodiments the nucleic acid is other than the entire retroviral genome of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, e.g., it is at least 1 nucleotide longer, or at least 1 nucleotide shorter, or differs in sequence at at least one position. E.g., the nucleic acid is a fragment of the sequence of SEQ ID NO:1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, or it includes sequence additional to that of SEQ ID NO:1, or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement.

20 In preferred embodiments: the sequence of the nucleic acid differs from the corresponding sequence of SEQ ID NO: 1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, by 1, 2, 3, 4, or 5 base pairs; the sequence of the nucleic acid differs from the corresponding sequence of SEQ ID NO: 1 or its complement, SEQ ID NO:2 or its complement, or SEQ ID NO:3 or its complement, by at least 1, 2, 3, 4, or 5 base pairs but less than 6, 7, 8, 9, or 10 base pairs.

25 In other preferred embodiments: the nucleic acid is at least 10, more preferably at least 15, more preferably at least 20, most preferably at least 25, 30, 50, 100, 1000, 2000, 30 4000, 6000, or 8060 nucleotides in length; the nucleic acid is less than 15, more preferably less than 20, most preferably less than 25, 30, 50, 100, 1000, 2000, 4000, 6000, or 8060 nucleotides in length.

Equivalents

35 Those skilled in the art will be able to recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures described herein. Such equivalents are considered to be within the scope of this invention and are covered by the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Jay A. Fishman

(ii) TITLE OF INVENTION: MOLECULAR SEQUENCE OF SWINE RETROVIRUS
AND METHODS OF USE

10

(iii) NUMBER OF SEQUENCES: 74

(iv) CORRESPONDENCE ADDRESS:

15

(A) ADDRESSEE: LAHIVE & COCKFIELD, LLP
(B) STREET: 60 State Street
(C) CITY: Boston
(D) STATE: Massachusetts
(E) COUNTRY: USA
(F) ZIP: 02109-1875

20

(v) COMPUTER READABLE FORM:

25

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

30

(A) APPLICATION NUMBER:
(B) FILING DATE:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/572,645
(B) FILING DATE: 14-DEC-1995

35

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Louis Myers
(B) REGISTRATION NUMBER: 35,965
(C) REFERENCE/DOCKET NUMBER: MGP-038CP

40

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (617)227-7400
(B) TELEFAX: (617)227-5941

45

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

50

(A) LENGTH: 8060 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CTCGAGACTC GGTGGAAGGG CCCTTATCTC GTACTTTTGA CCACACCAAC GGCTGTGAAA 60

GTCGAAGGAA TCTCCACCTG GATCCATGCA TCCCACGTTA AGCCGGCGCC ACCTCCCGAT 120
TCGGGGTGGA AAGCCGAAAA GACTGAAAAT CCCCTTAAGC TTCGCCTCCA TCGCGTGGTT 180
5 CCTTACTCTG TCAATAACCT CTCAGACTAA TGGTATGCGC ATAGGAGACA GCCTGAACTC 240
CCATAAACCC TTATCTCTCA CCTGGTTAAT TACTGACTCC GGCACAGGTA TTAATATCAA 300
10 CAACACTCAA GGGGAGGCTC CTTTAGGAAC CTGGTGGCCT GATCTATACG TTTGCCTCAG 360
ATCAGTTATT CCTAGTCTGA CCTCACCCCC AGATATCCTC CATGCTCACG GATTTTATGT 420
TTGCCCAGGA CCACCAAATA ATGGAAAACA TTGCGGAAAT CCCAGAGATT TCTTTTGTA 480
15 ACAATGGAAC TGTGTAACCT CTAATGATGG ATATTGGA 540
TAGGGTAAGT TTTTCTTATG TCAACACCTA TACCAGCTCT GGACAATTTA ATTACCTGAC 600
CTGGATTAGA ACTGGAAGCC CCAAGTGCTC TCCTTCAGAC CTAGATTACC TAAAAATAAG 660
20 TTTCACTGAG AAAGGAAAAC AAGAAAATAT CCTAAATGG GTAAATGGTA TGTCTTGGGG 720
AATGGTATAT TATGGAGGCT CGGGTAAACA ACCAGGCTCC ATTCTAACTA TTCGCCTCAA 780
25 AATAAACCAG CTGGAGCCTC CAATGGCTAT AGGACCAAAT ACGGTCTTGA CGGGTCAAAG 840
ACCCCAACC CAAGGACCAG GACCATCCTC TAACATAACT TCTGGATCAG ACCCACTGA 900
GTCTAGCAGC ACGACTAAAA TGGGGGCAAA ACTTTTTAGC CTCATCCAGG GAGCTTTTCA 960
30 AGCTCTTAAC TCCACGACTC CAGAGGCTAC CTCTTCTTGT TGGCTATGCT TAGCTTTGGG 1020
CCCACCTTAC TATGAAGGAA TGGCTAGAAG AGGGAAATTC AATGTGACAA AAGAACATAG 1080
35 AGACCAATGC ACATGGGGAT CCCAAAATAA GCTTACCCTT ACTGAGGTTT CTGGAAAAGG 1140
CACCTGCATA GGAAAGGTTT CCCCATCCCA CCAACACCTT TGTAACCACA CTGAAGCCTT 1200
TAATCAAACC TCTGAAAGTC AATATCTGGT ACCTGGTTAT GACAGGTGGT GGGCATGTAA 1260
40 TACTGGATTA ACCCCTTGTG TTTCCACCTT GGTTTTTAAC CAACTAAAG ATTTTTCAT 1320
TATGGTCCAA ATTGTTCCCC GAGTGATTA CTATCCCGAA AAAGCAATCC TTGATGAATA 1380
45 TGACTACAGA AATCATCGAC AAAAGAGAGA ACCCATATCT CTGACACTTG CTGTGATGCT 1440
CGGACTTGGA GTGGCAGCAG GTGTAGGAAC AGGAACAGCT GCCCTGGTCA CGGGACCACA 1500
GCAGCTAGAA ACAGGACTTA GTAACCTACA TCGAATTGTA ACAGAAGATC TCCAAGCCCT 1560
50 AGAAAAATCT GTCAGTAACC TGGAGGAATC CCTAACCTCC TTATCTGAAC TAGTCCTACA 1620
GAATAGAAGA GGGTTAGATT TATTATTTCT AAAAGAAGGA GGATTATGTG TAGCCTTGAA 1680
55 GGAGGAATGC TGTTTTTATG TGGATCATTC AGGGGCCATC AGAGACTCCA TGAACAACT 1740
TAGAGAAAGG TTGGAGAAGC GTCGAAGGGA AAAGGAACT ACTCAAGGGT GGT TTGAGGG 1800

ATGGTTCAAC AGGTCTCCTT GGTGGCTAC CCTACTTTCT GCTTTAACAG GACCCTTAAT 1860
5 AGTCCTCCTC CTGTTACTCA CAGTTGGGCC ATGTATTATT AACAAAGTTAA TTGCCTTCAT 1920
TAGAGAACGA ATAAGTGCAG TCCAGATCAT GGTACTTAGA CAACAGTACC AAAGCCCGTC 1980
TAGCAGGGAA GCTGGCCGCT AGCTCTACCA GTTCTAAGAT TAGAACTATT AACAAAGAGAA 2040
10 GAAGTGGGGA ATGAAAGGAT GAAAATACAA CCTAAGCTAA TGAGAAGCTT AAAATTGTTC 2100
TGAATTCCAG AGTTTGTTC TTATAGGTAA AAGATTAGGT TTTTGTCTGT TTTAAATAT 2160
GCGGAAGTAA AATAGGCCCT GAGTACATGT CTCTAGGCAT GAAACTTCTT GAACTATTT 2220
15 GAGATAACAA GAAAAGGGAG TTTCTAACTG CTTGTTTAGC TTCTGTAAAA CTGGTTGCGC 2280
CATAAAGATG TTGAAATGTT GATACACATA TCTTGGTGAC AACATGTCTC CCCACCCCG 2340
20 AAACATGCGC AAATGTGTAA CTCTAAAACA ATTTAAATTA ATTGGTCCAC GAAGCGCGGG 2400
CTCTCGAAGT TTTAAATTGA CTGGTTTGTG ATATTTTGAA ATGATTGGTT TGTAAGCGC 2460
GGGCTTTGCT GTGAACCCCA TAAAAGCTGT CCCGACTCCA CACTCGGGGC CGCAGTCCTC 2520
25 TACCCCTGCG TGGTGACGA CTGTGGGCCC CAGCGCGCTT GGAATAAAAA TCCTCTTGCT 2580
GTTTGATCA AGACCGCTTC TCGTGAGTGA TTAAGGGGAG TCGCCTTTTC CGAGCCTGGA 2640
30 GGTTCTTTTT GCTGGTCTTA CATTTGGGGG CTCGTCCGGG ATCTGTCGCG GCCACCCCTA 2700
ACACCCGAGA ACCGACTTGG AGGTAAAAAG GATCCTCTTT TTAACGTGTA TGCATGTACC 2760
GGCCGGCGTC TCTGTCTGA GTGTCTGTTT TCAGTGGTGC GCGCTTTCGG TTTGCAGCTG 2820
35 TCCTCTCAGG CCGTAAGGGC TGGGGGACTG TGATCAGCAG ACGTGCTAGG AGGATCACAG 2880
GCTGCTGCCC TGGGGGACGC CCCGGGAGGT GAGGAGAGCC AGGGACGCCT GGTGGTCTCC 2940
40 TACTGTCGGT CAGAGGACCG AATTCTGTTG CTGAAGCGAA AGCTTCCCCC TCCGCGACCG 3000
TCCGACTCTT TTGCCTGCTT GTGGAATACG TGGACGGGTC ACGTGTGTCT GGATCTGTTG 3060
GTTTCTGTTT TGTGTGTCTT TGTCTGTGT GTCTTGTCT ACAGTTTTAA TATGGGACAG 3120
45 ACGGTGACGA CCCCTCTTAG TTTGACTCTC GACCATTGGA CTGAAGTTAA ATCCAGGGCT 3180
CATAAATTGT CAGTTCAGGT TAAGAAGGGA CCTTGGCAGA CTTTCTGTGT CTCTGAATGG 3240
50 CCGACATTCG ATGTTGGATG GCCATCAGAG GGGACCTTTA ATTCTGAGAT TATCCTGGCT 3300
GTTAAAGCAA TTATTTTCA GACTGGACCC GGCTCTCATC CCGATCAGGA GCCCTATATC 3360
CTTACGTGGC AAGATTGGC AGAGGATCCT CCGCCATGGG TTAAACCATG GCTGAATAAG 3420
55 CCAAGAAAGC CAGGTCCCCG AATTCTGGCT CTTGGAGAGA AAAACAAACA CTCGGCTGAA 3480

AAAGTCAAGC CCTCTCCTCA TATCTACCCC GAGATTGAGG AACCACCGGC TTGGCCGGAA 3540
CCCCAATCTG TTCCCCCACC CCCTTATCTG GCACAGGGTG CCGCGAGGGG ACCCTTTGCC 3600
5 CCTCCTGGAG CTCCGGCGGT GGAGGGACCT TCTGCAGGGA CTCGGAGCCG GAGGGGCGCC 3660
ACCCCGGAGC GGACAGACGA GATCGCGACA TTACCGCTGC GCACGTACGG CCCTCCCACA 3720
10 CCGGGGGGCC AATTGCAGCC CCTCCAGTAT TGGCCCTTTT CTTCTGCAGA TCTCTATAAT 3780
TGGAAACTA ACCATCCCCC TTTCTCGGAG GATCCCCAAC GCCTCACGGG GTTGGTGGAG 3840
TCCCTTATGT TCTCTACCA GCCTACTTGG GATGATTGTC AACAGCTGCT GCAGACACTC 3900
15 TTCACAACCG AGGAGCGAGA GAGAATTCTA TTAGAGGCTA GAAAAATGT TCCTGGGGCC 3960
GACGGGCGAC CCACGCGGTT GCAAAATGAG ATTGACATGG GATTTCCCTT AACTCGCCCC 4020
GGTTGGGACT ACAACACGGC TGAAGGTAGG GAGAGCTTGA AAATCTATCG CCAGGCTCTG 4080
20 GTGGCGGGTC TCCGGGGCGC CTCAAGACGG CCCACTAATT TGGCTAAGGT AAGAGAAGTG 4140
ATGCAGGGAC CGAATGAACC CCCCTCTGTT TTTCTTGAGA GGCTCTTGGA AGCCTTCAGG 4200
25 CGGTACACCC CTTTTGATCC CACCTCAGAG GCCCAAAAAG CCTCAGTGGC TTTGGCCTTT 4260
ATAGGACAGT CAGCCTTGGA TATTAGAAAG AAGCTTCAGA GACTGGAAGG GTTACAGGAG 4320
GCTGAGTTAC GTGATCTAGT GAAGGAGGCA GAGAAAGTAT ATTACAAAAG GGAGACAGAA 4380
30 GAAGAAAGGG AACAAAGAAA AGAGAGAGAA AGAGAGGAAA GGGAGGAAAG ACGTAATAAA 4440
CGGCAAGAGA AGAATTTGAC TAAGATCTTG GCTGCAGTGG TTGAAGGGAA AAGCAATACG 4500
35 GAAAGAGAGA GAGATTTTAG GAAAATTAGG TCAGGCCCTA GACAGTCAGG GAACCTGGGC 4560
AATAGGACCC CACTCGACAA GGACCAATGT GCATATTGTA AAGAAAGAGG AACTGGGCA 4620
AGGAACTGCC CCAAGAAGGG AAACAAAGGA CCAAGGATCC TAGCTCTAGA AGAAGATAAA 4680
40 GATTAGGGGA GACGGGGTTC GGACCCCTC CCCGAGCCCA GGGTAACTTT GAAGGTGGAG 4740
GGGCAACCAG TTGAGTTCCT GGTGATACC GGAGCGAAAC ATTCACTGCT ACTACAGCCA 4800
45 TTAGGAAAAC TAAAAGATAA AAAATCCTGG GTGATGGGTG CACAGGGCAA CAACAGTATC 4860
CATGGACTAC CCGAAGACAG TTGACTTGGG AGTGGGACGG GTAACCCACT CGTTTCTGGT 4920
CATACTGAG TGCCCAGCAC CCCTCTTAGG TAGAGACTTA TTGACCAAGA TGGGAGCACA 4980
50 AATTTCTTTT GAACAAGGGA AACCAGAAGT GTCTGCAAT AACAAACCTA TCACTGTGTT 5040
GACCCTCAA TTAGATGACG AATATCGACT ATACTCTCCC CTAGTAAAGC CTGATCAAAA 5100
55 TATACAATTC TGGTTGGAAC AGTTTCCCA AGCCTGGGCA GAAACCGCAG GGATGGGTTT 5160
GGCAAAGCAA GTTCCCCAC AAGTTATTCA ACTGAAGGCC AGTGCCACAC CAGTGTCACT 5220

CAGACAGTAC CCCTTGAGTA AAGAAGCTCA AGAAGGAATT CGGCCGCATG TCCAAAGATT 5280
AATCCAACAG GGCATCCTAG TTCCTGTCCA ATCTCCCTGG AATACTCCCC TGCTACCGGT 5340
5 TAGAAAGCCT GGGACTAATG ACTATCGACC AGTACAGGAC TTGAGAGAGG TCAATAAACG 5400
GGTGCAGGAT ATACACCCAA CAGTCCCGAA CCCTTATAAC CTCTTGTTGT CTCTCCCACC 5460
10 CCAACGGAGC TGGTATACAG TATTGGACTT AAAGGATGCC TTCTTCTGCC TGAGATTACA 5520
CCCCACTAGC CAACCACTTT TTGCCTTCGA ATGGAGAGAT CCAGGTACGG GAAGAACCGG 5580
GCAGCTCACC TGGACCCGAC TGCCCCAAGG GTTCAAGAAC TCCCCGACCA TCTTTGACGA 5640
15 AGCCCTACAC AGAGACCTGG CCAACTTCAG GATCCAACAC CCTCAGGTGA CCCTCCTCCA 5700
GTACGTGGAT GACCTGCTTC TGGCGGGAGC CACCAAACAG GACTGCTTAG AAGGCACGAA 5760
20 GGCCTACTG CTGGAATTGT CTGACCTAGG CTACAGAGCC TCTGCTAAGA AGGCCCAGAT 5820
TTGCAGGAGA GAGGTAACAT ACTTGGGGTA CAGTTTACGG GACGGGCAGC GATGGCTGAC 5880
GGAGGCACGG AAGAAAACCTG TAGTCCAGAT ACCGGCCCCA ACCACAGCCA AACAAATGAG 5940
25 AGAGTTTTTG GGGACAGCTG GATTTTGCAG ACTGTGGATC CCGGGGTTTG CGACCTTAGC 6000
AGCCCCACTC TACCCGCTAA CCAAAGAAAA AGGGGAATTC TCCTGGGCTC CTGAGCACCA 6060
30 GAAGGCATTT GATGCTATCA AAAAGGCCCT GCTGAGCGCA CCTGCTCTGG CCCTCCCTGA 6120
CGTAACTAAA CCCTTTACCC TTTATGTGGA TGAGCGTAAG GGAGTAGCCC GGGGAGTTTT 6180
AACCCAAACC CTAGGACCAT GGAGAAGACC TGTCGCCTAC CTGTCAAAGA AGCTCGATCC 6240
35 TGTAGCCAGT GGTGGGCCA TATGCCTGAA GGCTATCGCA GCTGTGGCCA TACTGGTCAA 6300
GGACGCTGAC AAATTGACTT TGGGACAAGA ATATAACTGT AATAGCCCCC CATGCATTGG 6360
40 AGAACATCGT TCGGCAGCCC CCAGACCGAT GGATGACCAA CGCCCGCATG ACCCACTATC 6420
AAAGCCTGCT TCTCACAGAG AGGGTCACGT TCGCTCCACC AACCGCTCTC AACCTGCCA 6480
CTCTTCTGCC TGAAGAGACT GATGAACCAG TGAATCATGA TTGCCATCAA CTATTGATTG 6540
45 AGGAGACTGG GGTCCGCAAG GACCTTACAG ACATACCGCT GACTGGAGAA GTGCTAACCT 6600
GGTTCCTGA CGGAAGCAGC TATGTGGTGG AAGGTAAGAG GATGGCTGGG GCGGCGGTGG 6660
50 TGGACGGGAC CCGCACGATC TGGGCCAGCA GCCTGCCGGG AGGAACTTCA GCACAAAAGG 6720
CTGAGCTCAT GGCCCTCACG CAAGCTTTGC GGCTGGCCGA AGGGAAATCC ATAAACATTT 6780
ATACGGACAG CAGGTATGCC TTTGCGACTG CACACGTACA TGGGGCCATC TATAAACAAA 6840
55 GGGGGTTGCT TACCTCAGCA GGGAGGGAAA TAAAGAACAA AGAGGAAATT CTAAGCCTAT 6900

TAGAAGCCGT ACATTTACCA AAAAGGCTAG CTATTATACA CTGTCCTGGA CATCAGAAAG 6960
CTAAAGATCT CATATCCAGA GGAAACCAGA TGGCTGACCG GGTGCGCAAG CAGGCAGCCC 7020
5 AGGGTGTAA CCTTCTGCCT ATAATAGAAA TGCCCAAAGC CCCAGAACCC AGACGACAGT 7080
ACACCCTAGA AGACTGGCAA GAGATAAAAA AGATAGACCA TTCTCTGAGA CTCCGGAAGG 7140
10 GACCTGCTAT ACCTCAGATG GGAAGGAAAT CCTGCCCCAC AAAGAAGGGT TAGAATATGT 7200
CCAACAAGAT ACATCGTCTA ACCCACCTAG GAACTAAACA CCTGCAGCAG TTGGTCAGAA 7260
CATCCCCTTA TCATGTTCTG AGGCTACCAG GAGTGGCTGA CTCGGTGGTC AAACATTGTG 7320
15 TGCCCTGCCA GCTGGTTAAT GCTAATCCTT CCAGAATGCC TCCAGGAAG AGACTAAGGG 7380
GAAGCCACCC AGGCGCTCAC TGGGAAGTGG ACTTCACTGA GGTAAAGCCG GCTAAATATG 7440
GAAACAAATA CCTATTGGTT TTTGTAGACA CCTTTTCAGG ATGGGTAGAG GCTTATCCTA 7500
20 CTAAGAAAGA GACTTCAACC GTGGTAGCTA AAAAAATACT GGAAGAAATT TTTCCAAGAT 7560
TTGGAATACC TAAGGTAATA GGGTCAGACA ATGGTCCAGC TTTTGTGTC CAGGTAAGTC 7620
25 AGGGACTGGC CAAGATATTG GGGATTGATT GGAACTGCA TTGTGCATAC AGACCCCAAA 7680
GCTCAGGACA GGTAGAGAGG ATGAATAGAA CCATTAAAGA GACCCTTACT AAATTGACCG 7740
CGGAGACTGG CGTTAATGAT TGGATAGCTC TCCTGCCCTT TGTGCTTTTT AGGGTTAGGA 7800
30 ACACCCCTGG ACAGTTTGGG CTGACCCCTT ATGAATTACT CTACGGGGGA CCCCCCCAT 7860
TGGTAGAAAT TGCTTCTGTA CATAGTGCTG ATGTGCTGCT TTCCCAGCCT TTGTTCTCTA 7920
35 GGCTCAAGGC ACTTGAGTGG GTGAGACAAC GAGCGTGGAG GCAACTCCGG GAGGCCTACT 7980
CAGGAGGAGG AGACTTGCAAG ATCCACATC GTTTCCAAGT GGGAGATTCA GTCTACGTTA 8040
GACGCCACCG TGCAGGAAAC 8060

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 7333 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CTACCCCTGC GTGGTGTACG ACTGTGGGCC CCAGCGCGCT TGAATAAAAA ATCCTCTTGC 60
55 TGTTTGCATC AAGACCGCTT CTTGTGAGTG ATTTGGGGTG TCGCCTCTTC CGAGCCCGGA 120
CGAGGGGGAT TGTTCTTTTA CTGGCCTTTC ATTTGGTGCG TTGGCCGGGA AATCCTGCGA 180

CCACCCCTTA CACCCGAGAA CCGACTTGGA GGTAAAGGGA TCCCCTTTGG AACATATGTG 240
5 TGTGTCGGCC GGCGTCTCTG TTCTGAGTGT CTGTTTTTCGG TGATGCGCGC TTTCGGTTTTG 300
CAGCTGTCCT CTCAGACCGT AAGGACTGGA GGACTGTGAT CAGCAGACGT GCTAGGAGGA 360
TCACAGGCTG CCACCCCTGGG GGACGCCCCG GGAGGTGGGG AGAGCCAGGG ACGCCTGGTG 420
10 GTCTCCTACT GTCGGTCAGA GGACCGAGTT CTGTTGTTGA AGCGAAAGCT TCCCCCTCCG 480
CGGCCCTCCG ACTCTTTTGC CTGCTTGTGG AAGACGCGGA CGGGTCGCGT GTGTCTGGAT 540
15 CTGTTGGTTT CTGTTTCGTG TGTCTTTGTC TTGTGCGTCC TTGTCTACAG TTTTAATATG 600
GGACAGACAG TGA CTACCCC CCTTAGTTG ACTCTCGACC ATTGGACTGA AGTTAGATCC 660
AGGGCTCATA ATTTGTCAGT TCAGGTTAAG AAGGGACCTT GGCAGACTTT CTGTGCCTCT 720
20 GAATGGCCAA CATTCGATGT TGGATGGCCA TCAGAGGGGA CCTTTAATTC TGAAATTATC 780
CTGGCTGTTA AGGCAATCAT TTTTCAGACT GGACCCGGCT CTCATCCTGA TCAGGAGCCC 840
25 TATATCCTTA CGTGGCAAGA TTTGGCAGAA GATCCTCCGC CATGGGTAA ACCATGGCTA 900
AATAAACCAA GAAAGCCAGG TCCCCGAATC CTGGCTCTTG GAGAGAAAAA CAAACACTCG 960
GCCGAAAAAG TCGAGCCCTC TCCTCGTATC TACCCCGAGA TCGAGGAGCC GCCGACTTGG 1020
30 CCGGAACCCC AACCTGTTCC CCCACCCCT TATCCAGCAC AGGGTGCTGT GAGGGGACCC 1080
TCTGCCCTC CTGGAGCTCC GGTGGTGGAG GGACCTGCTG CCGGGACTCG GAGCCGGAGA 1140
GGCGCCACCC CGGAGCGGAC AGACGAGATC GCGATATTAC CGCTGCGCAC CTATGGCCCT 1200
35 CCCATGCCAG GGGGCCAATT GCAGCCCTC CAGTATTGGC CCTTTTCTTC TGCAGATCTC 1260
TATAATTGGA AAATAACCA TCCCCCTTC TCGGAGGATC CCCAACGCCT CACGGGGTTG 1320
40 GTGGAGTCCC TTATGTTCTC TCACCAGCCT ACTTGGGATG ATTGTCAACA GCTGCTGCAG 1380
ACACTCTTCA CAACCGAGGA GCGAGAGAGA ATTCTGTTAG AGGCTAAAAA AAATGTTTCCT 1440
GGGGCCGACG GGCGACCCAC GCAGTTGCAA AATGAGATTG ACATGGGATT TCCCTTGACT 1500
45 CGCCCCGGTT GGGACTACAA CACGGCTGAA GGTAGGGAGA GCTTGAAAAT CTATCGCCAG 1560
GCTCTGGTGG CGGGTCTCCG GGGCGCCTCA AGACGGCCCA CTAATTTGGC TAAGGTAAGA 1620
50 GAGGTGATGC AGGGACCGAA CGAACCTCCC TCGGTATTTT TTGAGAGGCT CATGGAAGCC 1680
TTCAGGCGGT TCACCCCTT TGATCCTACC TCAGAGGCC AGAAAGCCTC AGTGGCCCTG 1740
55 GCCTTCATTG GGCAGTCGGC TCTGGATATC AGGAAGAAAC TTCAGAGACT GGAAGGGTTA 1800
CAGGAGGCTG AGTTACGTGA TCTAGTGAGA GAGGCAGAGA AGGTGTATTA CAGAAGGGAG 1860

| | | | | | | | |
|----|-------------|------------|------------|------------|-------------|------------|------|
| | ACAGAAGAGG | AGAAGGAACA | GAGAAAAGAA | AAGGAGAGAG | AAGAAAGGGA | GGAAAGACGT | 1920 |
| | GATAGACGGC | AAGAGAAGAA | TTTGACTAAG | ATCTTGGCCG | CAGTGGTTGA | AGGGAAGAGC | 1980 |
| 5 | AGCAGGGAGA | GAGAGAGAGA | TTTTAGGAAA | ATTAGGTCAG | GCCCTAGACA | GTCAGGGAAC | 2040 |
| | CTGGGCAATA | GGACCCCACT | CGACAAGGAC | CAGTGTGCGT | ATTGTAAAGA | AAAAGGACAC | 2100 |
| | TGGGCAAGGA | ACTGCCCCAA | GAAGGGAAAC | AAAGGACCGA | AGGTCCCTAGC | TCTAGAAGAA | 2160 |
| 10 | GATAAAGATT | AGGGGAGACG | GGGTTCGGAC | CCCCTCCCCG | AGCCCAGGGT | AACTTTGAAG | 2220 |
| | GTGGAGGGGC | AACCAGTTGA | GTTCTGTTT | GATACCGGAG | CGGAGCATTC | AGTGCTGCTA | 2280 |
| 15 | CAACCATTAG | GAAAACATAA | AGAAAAAATA | TCCTGGGTGA | TGGGTGCCAC | AGGGCAACGG | 2340 |
| | CAGTATCCAT | GGACTACCCG | AAGAACCGTT | GACTTGGGAG | TGGGACGGGT | AACCCACTCG | 2400 |
| | TTTCTGGTCA | TCCCTGAGTG | CCCAGTACCC | CTTCTAGGTA | GAGACTTACT | GACCAAGATG | 2460 |
| 20 | GGAGCTCAAA | TTTCTTTTGA | ACAAGGAAGA | CCAGAAGTGT | CTGTGAATAA | CAAACCCATC | 2520 |
| | ACTGTGTTGA | CCCTCCAATT | AGATGATGAA | TATCGACTAT | ATTCTCCCCA | AGTAAAGCCT | 2580 |
| 25 | GATCAAGATA | TACAGTCCTG | GTTGGAGCAG | TTTCCCCAAG | CCTGGGCAGA | AACCGCAGGG | 2640 |
| | ATGGGTTTGG | CAAAGCAAGT | TCCCCCACAG | GTTATTCAAC | TGAAGGCCAG | TGCTACACCA | 2700 |
| | GTATCAGTCA | GACAGTACCC | CTTGAGTAGA | GAGGCTCGAG | AAGGAATTTG | GCCGCATGTT | 2760 |
| 30 | CAAAGATTAA | TCCAACAGGG | CATCCTAGTT | CCTGTCCAAT | CCCCTTGGAA | TACTCCCCTG | 2820 |
| | CTACCGGTTA | GGAAGCCTGG | GACCAATGAT | TATCGACCAG | TACAGGACTT | GAGAGAGGTC | 2880 |
| 35 | AATAAAAGGG | TGCAGGACAT | ACACCCAACG | GTCCCGAACC | CTTATAACCT | CTTGAGCGCC | 2940 |
| | CTCCCGCCTG | AACGGAAGTG | GTACACAGTA | TTGGACTTAA | AAGATGCCTT | CTTCTGCCTG | 3000 |
| | AGATTACACC | CCACTAGCCA | ACCACTTTTT | ACCTTCGAAT | GGAGAGATCC | AGGTACGGGA | 3060 |
| 40 | AGAACCGGGC | AGCTCACCTG | GACCCGACTG | CCCCAAGGGT | TCAAGAACTC | CCCGACCATC | 3120 |
| | TTTGACGAAG | CCCTACACAG | GGACCTGGCC | AACTTCAGGA | TCCAACACCC | TCAGGTGACC | 3180 |
| 45 | CTCCTCCAGT | ACGTGGATGA | CCTGCTTCTG | GCGGGAGCCA | CCAAACAGGA | CTGCTTAGAA | 3240 |
| | GGTACGAAGG | CACTACTGCT | GGAATTGTCT | GACCTAGGCT | ACAGAGCCTC | TGCTAAGAAG | 3300 |
| | GCCCAGATTT | GCAGGAGAGA | GGTAACATAC | TTGGGGTACA | GTTTGCGGGG | CGGGCAGCGA | 3360 |
| 50 | TGGCTGACGG | AGGCACGGAA | GAAAACTGTA | GTCCAGATAC | CGGCCCAAC | CACAGCCAAA | 3420 |
| | CAAGTGAGAG | AGTTTTTGGG | GACAGCTGGA | TTTTGCAGAC | TGTGGATCCC | GGGGTTTGCG | 3480 |
| 55 | ACCTTAGCAG | CCCCACTCTA | CCCGCTAACC | AAAGAAAAAG | GGGGTTGCTT | ACCTCAGCAG | 3540 |
| | GGAGGGAAAAT | AAAGAACAAA | GAGGAAATTC | TAAGCCTATT | AGAAGCCTTA | CATTTGCCAA | 3600 |

| | | | | | | | |
|----|------------|------------|------------|------------|------------|------------|------|
| | AAAGGCTAGC | TATTATACAC | TGTCCTGGAC | ATCAGAAAGC | CAAAGATCTC | ATATCTAGAG | 3660 |
| 5 | GGAACCAGAT | GGCTGACCGG | GTTGCCAAGC | AGGCAGCCCA | GGCTGTTAAC | CTTCTGCCTA | 3720 |
| | TAATAGAAAC | GCCCAAAGCC | CCAGAACCCA | GACGACAGTA | CACCCTAGAA | GACTGGCAAG | 3780 |
| | AGATAAAAAA | GATAGACCAG | TTCTCTGAGA | CTCCGGAGGG | GACCTGCTAT | ACCTCATATG | 3840 |
| 10 | GGAAGGAAAT | CCTGCCCCAC | AAAGAAGGGT | TAGAATATGT | CCAACAGATA | CATCGTCTAA | 3900 |
| | CCCACCTAGG | AACTAAACAC | CTGCAGCAGT | TGGTCAGAAC | ATCCCCTTAT | CATGTTCTGA | 3960 |
| 15 | GGCTACCAGG | AGTGGCTGAC | TCGGTGGTCA | AACATTGTGT | GCCCTGCCAG | CTGGTTAATG | 4020 |
| | CTAATCCTTC | CAGAATACCT | CCAGGAAAGA | GACTAAGGGG | AAGCCACCCA | GGCGCTCACT | 4080 |
| | GGGAAGTGGA | CTTCACTGAG | GTAAAGCCGG | CTAAATACGG | AAACAAATAT | CTATTGGTTT | 4140 |
| 20 | TTGTAGACAC | CTTTTCAGGA | TGGGTAGAGG | CTTATCCTAC | TAAAAAAGAG | ACTTCAACCG | 4200 |
| | TGGTGGCTAA | GAAAATACTG | GAGGAAATTT | TTCCAAGATT | TGGAATACCT | AAGGTAATAG | 4260 |
| 25 | GGTCAGACAA | TGGTCCAGCT | TTCGTTGCCC | AGGTAAGTCA | GGGACTGGCC | AAGATATTGG | 4320 |
| | GGATTGATTG | AAAACGTCAT | TGTGCATACA | GACCCCAAAG | CTCAGGACAG | GTAGAGAGGA | 4380 |
| | TGAATAGAAC | CATTAAAGAG | ACCCTTACCA | AATTGACCAC | AGAGACTGGC | ATTAATGATT | 4440 |
| 30 | GGATGGCTCT | CCTGCCCTTT | GTGCTTTTTA | GGGTGAGGAA | CACCCCTGGA | CAGTTTGGGC | 4500 |
| | TGACCCCTTA | TAAATTGCTC | TACGGGGGAC | CCCCCCCCTT | GGCAGAAATT | GCCTTTGCAC | 4560 |
| 35 | ATAGTGCTGA | TGTGCTGCTT | TCCCAGCCTT | TGTTCTCTAG | GCTCAAGGCG | CTCGAGTGGG | 4620 |
| | TGAGGCAGCG | AGCGTGGAAG | CAGCTCCGGG | AGGCCTACTC | AGGAGGAGAC | TTGCAAGTTC | 4680 |
| | CACATCGCTT | CCAAGTTGGA | GATTCACTCT | ATGTTAGACG | CCACCGTGCA | GGAAACCTCG | 4740 |
| 40 | AGACTCGGTA | GAAGGGACCT | TATCTCGTAC | TTTTGACCAC | ACCAACGGCT | GTGAAAGTCG | 4800 |
| | AAGGAATCCC | CTTAAGCTTC | GCCTCCATCG | CGTGGTTCCT | TACTCTGTCA | ATAACTCCTC | 4860 |
| 45 | AAGTTAATGG | TAAACGCCTT | GTGGACAGCC | CGAACTCCCA | TAAACCCTTA | TCTCTCACCT | 4920 |
| | GGTTACTTAC | TGACTCCGGT | ACAGGTATTA | ATATTAACAG | CACTCAAGGG | GAGGCTCCCT | 4980 |
| | TGGGGACCTG | GTGGCCTGAA | TTATATGTCT | GCCTTCGATC | AGTAATCCCT | GGTCTCAATG | 5040 |
| 50 | ACCAGGCCAC | ACCCCCCGAT | GTACTCCGTG | CTTACGGGTT | TTACGTTTGC | CCAGGACCCC | 5100 |
| | CAAATAATGA | AGAATATTGT | GGAAATCCTC | AGGATTTCTT | TTGCAAGCAA | TGGAGCTGCA | 5160 |
| 55 | TAACCTCTAA | TGATGGGAAT | TGGAAATGGC | CAGTCTCTCA | GCAAGACAGA | GTAAGTTACT | 5220 |
| | CTTTTGTTAA | CAATCCTACC | AGTTATAATC | AATTTAATTA | TGGCCATGGG | AGATGGAAAG | 5280 |

| | | | | | | | |
|----|------------|-------------|------------|------------|------------|-------------|------|
| | ATTGGCAACA | GCGGGTACAA | AAAGATGTAC | GAAATAAGCA | AATAAGCTGT | CATTTCGTTAG | 5340 |
| | ACCTAGATTA | CTTAAAAATA | AGTTTCACTG | AAAAAGGAAA | ACAAGAAAAT | ATTCAAAAAGT | 5400 |
| 5 | GCGTAAATGG | TATATCTTGG | GGAATAGTGT | ACTATGGAGG | CTCTGGGAGA | AAGAAAGGAT | 5460 |
| | CTGTTCTGAC | TATTTCGCCTC | AGAATAGAAA | CTCAGATGGA | ACCTCCGGTT | GCTATAGGAC | 5520 |
| | CAAATAAGGG | TTTGCCGAA | CAAGGACCTC | CAATCCAAGA | ACAGAGGCCA | TCTCCTAACC | 5580 |
| 10 | CCTCTGATTA | CAATACAACC | TCTGGATCAG | TCCCCACTGA | GCCTAACATC | ACTATTAAAA | 5640 |
| | CAGGGGCGAA | ACTTTTTAGC | CTCATCCAGG | GAGCTTTTCA | AGCTCTTAAC | TCCACGACTC | 5700 |
| 15 | CAGAGGCTAC | CTCTTCTTGT | TGGCTTTGCT | TAGCTTCGGG | CCCACCTTAC | TATGAGGGAA | 5760 |
| | TGGCTAGAGG | AGGGAAATTC | AATGTGACAA | AGGAACATAG | AGACCAATGT | ACATGGGGAT | 5820 |
| | CCCAAAATAA | GCTTACCCTT | ACTGAGGTTT | CTGGAAAAGG | CACCTGCATA | GGGATGGTTC | 5880 |
| 20 | CCCCATCCCA | CCAACACCTT | TGTAACCACA | CTGAAGCCTT | TAATCGAACC | TCTGAGAGTC | 5940 |
| | AATATCTGGT | ACCTGGTTAT | GACAGGTGGT | GGGCATGTAA | TACTGGATTA | ACCCCTTGTTG | 6000 |
| 25 | TTTCCACCTT | GGTTTTCAAC | CAAATAAAG | ACTTTTGCGT | TATGGTCCAA | ATTGTCCCCC | 6060 |
| | GGGTGTACTA | CTATCCCGAA | AAAGCAGTCC | TTGATGAATA | TGACTATAGA | TATAATCGGC | 6120 |
| | CAAAAAGAGA | GCCCATATCC | CTGACACTAG | CTGTAATGCT | CGGATTGGGA | GTGGCTGCAG | 6180 |
| 30 | GCGTGGGAAC | AGGAACGGCT | GCCCTAATCA | CAGGACCGCA | ACAGCTGGAG | AAAGGACTTA | 6240 |
| | GTAACCTACA | TCGAATTGTA | ACGGAAGATC | TCCAAGCCCT | AGAAAAATCT | GTCAGTAACC | 6300 |
| 35 | TGGAGGAATC | CCTAACCTCC | TTATCTGAAG | TGGTTCTACA | GAACAGAAGG | GGGTTAGATC | 6360 |
| | TGTTATTCT | AAAAGAAGGA | GGGTTATGTG | TAGCCTTAAA | AGAGGAATGC | TGCTTCTATG | 6420 |
| | TAGATCACTC | AGGAGCCATC | AGAGACTCCA | TGAGCAAGCT | TAGAGAAAGG | TTAGAGAGGC | 6480 |
| 40 | GTCGAAGGGA | AAGAGAGGCT | GACCAGGGGT | GGTTTGAAGG | ATGGTTCAAC | AGGTCTCCTT | 6540 |
| | GGATGACCAC | CCTGCTTTCT | GCTCTGACGG | GGCCCCTAGT | AGTCCTGCTC | CTGTTACTTA | 6600 |
| 45 | CAGTTGGGCC | TTGCTTAATT | AATAGGTTTG | TTGCCTTTGT | TAGAGAACGA | GTGAGTGCAG | 6660 |
| | TCCAGATCAT | GGTACTTAGG | CAACAGTACC | AAGGCCTTCT | GAGCCAAGGA | GAAACTGACC | 6720 |
| | TCTAGCCTTC | CCAGTTCTAA | GATTAGAACT | ATTAACAAGA | CAAGAAGTGG | GGAATGAAAG | 6780 |
| 50 | GATGAAAATG | CAACCTAACC | CTCCCAGAAC | CCAGGAAGTT | AATAAAAAGC | TCTAAATGCC | 6840 |
| | CCCGAATTCC | AGACCTTGCT | GGCTGCCAGT | AAATAGGTAG | AAGGTCACAC | TTCTATTGT | 6900 |
| 55 | TCCAGGGCCT | GCTATCCTGG | CCTAAGTAAG | ATAACAGGAA | ATGAGTTGAC | TAATCGCTTA | 6960 |
| | TCTGGATTCT | GTAAAACTGA | CTGGCACCAT | AGAAGAATTG | ATTACACATT | GACAGCCCTA | 7020 |

GTGACCTATC TCAACTGCAA TCTGTCACTC TGCCCAGGAG CCCACGCAGA TGCGGACCTC 7080
CGGAGCTATT TAAAATGAT TGGTCCACGG AGCGCGGGCT CTCGATATTT TAAAATGATT 7140
5 GGTCCATGGA GCGCGGGCTC TCGATATTTT AAAATGATTG GTTTGTGACG CACAGGCTTT 7200
GTTGTGAACC CCATAAAGC TGTCCCATT CCGCACTCGG GGCCGCAGTC CTCTACCCCT 7260
10 GCGTGGTGTA CGACTGTGGG CCCAGCGCG CTTGGAATAA AAATCCTCTT GCTGTTTGCA 7320
TCAAAAAAAAA AAA 7333

(2) INFORMATION FOR SEQ ID NO:3:

15

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8132 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
20 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GCGTGGTGTA CGACTGTGGG CCCAGCGCG CTTGGAATAA AAATCCTCTT GCTGTTTGCA 60
30 TCAAGACCGC TTCTCGTGAG TGATTAAGGG GAGTCGCCTT TTCCGAGCCT GGAGGTTCTT 120
TTTGCTGGTC TTACATTTGG GGGCTCGTCC GGGATCTGTC GCGGCCACCC CTAACACCCG 180
35 AGAACCGACT TGGAGGTAAA AAGGATCCTC TTTTAAACGT GTATGCATGT ACCGGCCGGC 240
GTCTCTGTTT TGAGTGTCTG TTTTCAGTGG TGCGCGCTTT CGGTTTGAGC CTGTCTCTCTC 300
AGGCCGTAAG GGCTGGGGGA CTGTGATCAG CAGACGTGCT AGGAGGATCA CAGGCTGCTG 360
40 CCCTGGGGGA CGCCCCGGA GGTGAGGAGA GCCAGGGACG CCTGGTGGTC TCCTACTGTC 420
GGTCAGAGGA CCGAATTCTG TTGCTGAAGC GAAAGCTTCC CCCTCCGCGA CCGTCCGACT 480
45 CTTTTGCCTG CTTGTGGAAG ACGTGGACGG GTCACGTGTG TCTGGATCTG TTGGTTTCTG 540
TTTTGTGTGT CTTGTCTTG TGTGTCCTTG TCTACAGTTT TAATATGGGA CAGACGGTGA 600
CGACCCCTCT TAGTTTGA CTGACCATTT GGACTGAAGT TAAATCCAGG GCTCATAATT 660
50 TGTCAGTTCA GGTAAAGAAG GGACCTTGGC AGACTTTCTG TGTCTCTGAA TGGCCGACAT 720
TCGATGTTGG ATGGCCATCA GAGGGGACCT TTAATTCTGA GATTATCCTG GCTGTAAAG 780
55 CAGTTATTTT TCAGACTGGA CCCGGCTCTC ATCCCGATCA GGAGCCCTAT ATCCTTACGT 840
GGCAAGATTT GGCAGAGGAT CCTCCGCCAT GGGTTAAACC ATGGCTGAAT AAGCCAAGAA 900

| | | | | | | | |
|----|------------|------------|------------|------------|-------------|------------|------|
| | AGCCAGGTCC | CCGAATTCTG | GCTCTTGGAG | AGAAAAACAA | AACTTCGGCT | GAAAAAGTCA | 960 |
| | AGCCCTCTCC | TCATATCTAC | CCCGAGATTG | AGGAGCCACC | GGCTTGCCCG | GAACCCCAAT | 1020 |
| 5 | CTGTTCCCCC | ACCCCTTAT | CTGGCACAGG | GTGCCGCGAG | GGGACCCTTT | GCCCCCTCTG | 1080 |
| | GAGCTCCGGC | GGTGGAGGGA | CCTGCTGCAG | GGACTCGGAG | CCGGAGGGGC | GCCACCCCGG | 1140 |
| 10 | AGCGGACAGA | CGAGATCGCG | ACATTACCGC | TGCGCACGTA | CGGCCCTCCC | ACACCGGGGG | 1200 |
| | GCCAATTGCA | GCCCCCTCAG | TATTGGCCCT | TTTCTTCTGC | AGATCTCTAT | AATTGGAAAA | 1260 |
| | CTAACCATCC | CCCTTTCTCG | GAGGATCCCC | AACGCCTCAC | GGGGTTGGTG | GAGTCCCTTA | 1320 |
| 15 | TGTTCTCTCA | CCAGCCTACT | TGGGATGATT | GTCAACAGCT | GCTGCAGACA | CTCTTCACAA | 1380 |
| | CCGAGGAGCG | AGAGAGAATT | CTATTAGAGG | CTAGAAAAAA | TGTTCCCTGGG | GCCGACGGGC | 1440 |
| 20 | GACCCACGCG | GTTGCAAAAT | GAGATTGACA | TGGGATTTC | CTTAACTCGC | CCCGTTGGG | 1500 |
| | ACTACAACAC | GGCTGAAGGT | AGGGAGAGCT | TGAAAATCTA | TCGCCAGGCT | CTGGTGGCGG | 1560 |
| | GTCTCCGGGG | CGCCTCAAGA | CGGCCCACTA | ATTTGGCTAA | GGTAAGAGAA | GTGATGCAGG | 1620 |
| 25 | GACCGAATGA | ACCCCTCTCT | GTTTTTCTTG | AGAGGCTCTT | GGAAGCCTTC | AGGCGGTACA | 1680 |
| | CCCCTTTTGA | TCCCACCTCA | GAGGCCCAA | AAGCCTCAGT | GGCTTTGGCC | TTTATAGGAC | 1740 |
| 30 | AGTCAGCCTT | GGATATTAGA | AAGAAGCTTC | AGAGACTGGA | AGGGTTACAG | GAGGCTGAGT | 1800 |
| | TACGTGATCT | AGTGAAGGAG | GCAGAGAAAG | TATATTACAA | AAGGGAGACA | GAAGAAGAAA | 1860 |
| | GGGAACAAAG | AAAAGAGAGA | GAAAGAGAGG | AAAGGGAGGA | AAGACGTAAT | AAACGGCAAG | 1920 |
| 35 | AGAAGAATTT | GACTAAGATC | TTGGCTGCAG | TGGTTGAAGG | GAAAAGCAAT | ACGGAAAGAG | 1980 |
| | AGAGAGATTT | TAGGAAAATT | AGGTCAGGCC | CTAGACAGTC | AGGGAACCTG | GGCAATAGGA | 2040 |
| 40 | CCCCACTCGA | CAAGGACCAA | TGTGCATATT | GTAAAGAAAG | AGGACACTGG | GCAAGGAACT | 2100 |
| | GCCCCAAGAA | GGGAAACAAA | GGACCAAGGA | TCCTAGCTCT | AGAAGAAGAT | AAAGATTAGG | 2160 |
| | GGAGACGGGG | TTCGGACCCC | CTCCCCGAGC | CCAGGGTAAC | TTTGAAGGTG | GAGGGGCAAC | 2220 |
| 45 | CAGTTGAGTT | CCTGGTTGAT | ACCGGAGCGA | AACATTCACT | GCTACTACAG | CCATTAGGAA | 2280 |
| | AACTAAAAGA | TAAAAAATCC | TGGGTGATGG | GTGCCACAGG | GCAACAACAG | TATCCATGGA | 2340 |
| 50 | CTACCCGAAG | AACAGTTGAC | TTGGGAGTGG | GACGGGTAA | CCACTCGTTT | CTGGTCATAC | 2400 |
| | CTGAGTGCCC | AGCACCCCTC | TTAGGTAGAG | ACTTATTGAC | CAAGATGGGA | GCACAAATTT | 2460 |
| | CTTTTGAACA | AGGGAAACCA | GAAGTGTCTG | CAAATAACAA | ACCTATCACT | GTGTTGACCC | 2520 |
| 55 | TCCAATTAGA | TGACGAATAT | CGACTATACT | CTCCCTAGT | AAAGCCTGAT | CAAAATATAC | 2580 |

| | | | | | | | |
|----|-------------|-------------|------------|------------|------------|------------|------|
| | AATTCTGGTT | GGAACAGTTT | CCCCAAGCCT | GGGCAGAAAC | CGCAGGGATG | GGTTTGGCAA | 2640 |
| | AGCAAGTTCC | CCCACAAGTT | ATTCAACTGA | AGGCCAGTGC | CACACCAGTG | TCAGTCAGAC | 2700 |
| 5 | AGTACCCCTT | GAGTAAAGAA | GCTCAAGAAG | GAATTCGGCC | GCATGTCCAA | AGATTAATCC | 2760 |
| | AACAGGGCAT | CCTAGTTCCT | GTCCAATCTC | CCTGGAATAC | TCCCCTGCTA | CCGGTTAGAA | 2820 |
| | AGCCTGGGAC | TAATGACTAT | CGACCAGTAC | AGGACTTGAG | AGAGGTCAAT | AAACGGGTGC | 2880 |
| 10 | AGGATATACA | CCCAACAGTC | CCGAACCCTT | ATAACCTCTT | GTGTGCTCTC | CCACCCCAAC | 2940 |
| | GGAGCTGGTA | TACAGTATTG | GACTTAAAGG | ATGCCTTCTT | CTGCCTGAGA | TTACACCCCA | 3000 |
| 15 | CTAGCCAACC | ACTTTTGGCC | TTCGAATGGA | GAGATCCAGG | TACGGGAAGA | ACCGGGCAGC | 3060 |
| | TCACCTGGAC | CCGACTGCCC | CAAGGGTTCA | AGAACTCCCC | GACCATCTTT | GACGAAGCCC | 3120 |
| | TACACAGAGA | CCTGGCCAAC | TTCAGGATCC | AACACCCTCA | GGTGACCCTC | CTCCAGTACG | 3180 |
| 20 | TGGATGACCT | GCTTCTGGCG | GGAGCCACCA | AACAGGACTG | CTTAGAAGGC | ACGAAGGCAC | 3240 |
| | TACTGCTGGA | ATTGTCTGAC | CTAGGCTACA | GAGCCTCTGC | TAAGAAGGCC | CAGATTTGCA | 3300 |
| 25 | GGAGAGAGGT | AACATACTTG | GGGTACAGTT | TGCGGGACGG | GCAGCGATGG | CTGACGGAGG | 3360 |
| | CACGGAAGAA | AACTGTAGTC | CAGATACCGG | CCCCAACCAC | AGCCAAACAA | ATGAGAGAGT | 3420 |
| | TTTTGGGGAC | AGCTGGATTT | TGCAGACTGT | GGATCCCGGG | GTTTGCGACC | TTAGCAGCCC | 3480 |
| 30 | CACTCTACCC | GCTAACCAAA | GAAAAAGGGG | AATTCTCTCG | GGCTCCTGAG | CACCAGAAGG | 3540 |
| | CATTTGATGC | TATCAAAAAG | GCCCTGCTGA | GCGCACCTGC | TCTGGCCCTC | CCTGACGTAA | 3600 |
| 35 | CTAAACCCTT | TACCCTTTAT | GTGGATGAGC | GTAAGGGAGT | AGCCCGGGGA | GTTTTAACCC | 3660 |
| | AAACCCTAGG | ACCATGGAGA | AGACCTGTCT | CCTACCTGTC | AAAGAAGCTC | GATCCTGTAG | 3720 |
| | CCAGTG GTTG | GCCCATATGC | CTGAAGGCTA | TCGCAGCTGT | GGCCATACTG | GTCAAGGACG | 3780 |
| 40 | CTGACAAATT | GACTTTGGGA | CAGAAATATA | CTGTAATAGC | CCCCCATGCA | TTGGAGAACA | 3840 |
| | TCGTTTCGGCA | GCCCCCAGAC | CGATGGATGA | CCAACGCCCC | CATGACCCAC | TATCAAAGCC | 3900 |
| 45 | TGCTTCTCAC | AGAGAGGGTC | ACGTTCTGCT | CACCAGCCGC | TCTCAACCCT | GCCACTCTTC | 3960 |
| | TGCCTGAAGA | GA CTGATGAA | CCAGTGACTC | ATGATTGCCA | TCAACTATTG | ATTGAGGAGA | 4020 |
| | CTGGGGTCCG | CAAGGACCTT | ACAGACATAC | CGCTGACTGG | AGAAGTGCTA | ACCTGGTTCA | 4080 |
| 50 | CTGACGGAAG | CAGCTATGTG | GTGGAAGGTA | AGAGGATGGC | TGGGGCGGCG | GTGGTGGACG | 4140 |
| | GGACCCGCAC | GATCTGGGCC | AGCAGCCTGC | CGGAAGGAAC | TTCAGCACAA | AAGGCTGAGC | 4200 |
| 55 | TCATGGCCCT | CACGCAAGCT | TTGCGGCTGG | CCGAAGGGAA | ATCCATAAAC | ATTTATACGG | 4260 |
| | ACAGCAGGTA | TGCCTTTGCG | ACTGCACACG | TACATGGGGC | CATCTATAAA | CAAAGGGGGT | 4320 |

| | | | | | | | |
|----|------------|------------|------------|------------|------------|------------|------|
| | TGCTTACCTC | AGCAGGGAGG | GAAATAAAGA | ACAAAGAGGA | AATTCTAAGC | CTATTAGAAG | 4380 |
| | CCGTACATTT | ACCAAAAAGG | CTAGCTATTA | TACACTGTCC | TGGACATCAG | AAAGCTAAAG | 4440 |
| 5 | ATCTCATATC | CAGAGGAAAC | CAGATGGCTG | ACCGGGTTGC | CAAGCAGGCA | GCCCAGGGTG | 4500 |
| | TTAACCTTCT | GCCTATAATA | GAAATGCCCA | AAGCCCCAGA | ACCCAGACGA | CAGTACACCC | 4560 |
| 10 | TAGAAGACTG | GCAAGAGATA | AAAAAGATAG | ACCAGTTCTC | TGAGACTCCG | GAAGGGACCT | 4620 |
| | GCTATACCTC | AGATGGGAAG | GAAATCCTGC | CCCACAAAGA | AGGGTTAGAA | TATGTCCAAC | 4680 |
| | AGATACATCG | TCTAACCCAC | CTAGGAACTA | AACACCTGCA | GCAGTTGGTC | AGAACATCCC | 4740 |
| 15 | CTTATCATGT | TCTGAGGCTA | CCAGGAGTGG | CTGACTCGGT | GGTCAAACAT | TGTGTGCCCT | 4800 |
| | GCCAGCTGGT | TAATGCTAAT | CCTTCCAGAA | TGCCTCCAGG | GAAGAGACTA | AGGGGAAGCC | 4860 |
| 20 | ACCCAGGCGC | TCACTGGGAA | GTGGACTTCA | CTGAGGTAAA | GCCGGCTAAA | TACGGAAACA | 4920 |
| | AATACCTATT | GGTTTTTGTA | GACACCTTTT | CAGGATGGGT | AGAGGCTTAT | CCTACTAAGA | 4980 |
| | AAGAGACTTC | AACCGTGGTG | GCTAAAAAAA | TACTGGAAGA | AATTTTCCA | AGATTTGGAA | 5040 |
| 25 | TACCTAAGGT | AATAGGGTCA | GACAATGGTC | CAGCTTTTGT | TGCCCAGGTA | AGTCAGGGAC | 5100 |
| | TGGCCAAGAT | ATTGGGGATT | GATTGGAAAC | TGCATTGTGC | ATACAGACCC | CAAAGCTCAG | 5160 |
| 30 | GACAGGTAGA | GAGGATGAAT | AGAACCATTA | AAGAGACCCT | TACTAAATTG | ACCGCGGAGA | 5220 |
| | CTGGCGTTAA | TGATTGGATA | GCTCTCCTGC | CCTTTGTGCT | TTTtagGGTT | AGGAACACCC | 5280 |
| | CTGGACAGTT | TGGGCTGACC | CCCTATGAAT | TACTCTACGG | GGGACCCCCC | CCATTGGTAG | 5340 |
| 35 | AAATTGCTTC | TGTACATAGT | GCTGACGTGC | TGCTTTCCCA | GCCTTTGTTC | TCTAGGCTCA | 5400 |
| | AGGCACTTGA | GTGGGTGAGA | CAACGAGCGT | GGAGGCAACT | CCGGGAGGCC | TACTCAGGAG | 5460 |
| 40 | GAGGAGACTT | GCAGATCCCA | CATCGTTTCC | AAGTGGGAGA | TTCAGTCTAC | GTTAGACGCC | 5520 |
| | ACCGTGCAGG | AAACCTCGAG | ACTCGGTGGA | AGGGCCCTTA | TCTCGTACTT | TTGACCACAC | 5580 |
| | CAACGGCTGT | GAAAGTCGAA | GGAATCTCCA | CCTGGATCCA | TGCATCCCAC | GTTAAACCGG | 5640 |
| 45 | CGCCACCTCC | CGATTGCGGG | TGGAAAGCCG | AAAAGACTGA | AAATCCCCTT | AAGCTTCGCC | 5700 |
| | TCCATCGCGT | GGTTCCTTAC | TCTGTCAATA | ACCTCTCAGA | CTAATGGTAT | GCGCATAGGA | 5760 |
| 50 | GACAGCCTGA | ACTCCCATAA | ACCCTTATCT | CTCACCTGGT | TAATTACTGA | CTCCGGCACA | 5820 |
| | GGTATTAATA | TCAACAACAC | TCAAGGGGAG | GCTCCTTTAG | GAACCTGGTG | GCCTGATCTA | 5880 |
| | TACGTTTGCC | TCAGATCAGT | TATTCCTAGT | CTGACCTCAC | CCCCAGATAT | CCTCCAIGCT | 5940 |
| 55 | CACGGATTTT | ATGTTTGCCC | AGGACCACCA | AATAATGGAA | AACATTGCGG | AAATCCCAGA | 6000 |

| | | | | | | | |
|----|------------|------------|-------------|------------|------------|------------|------|
| | GATTTCTTTT | GTAAACAATG | GAAGTGTGTA | ACCTCTAATG | ATGGATATTG | GAAATGGCCA | 6060 |
| | ACCTCTCAGC | AGGATAGGGT | AAGTTTTTCT | TATGTCAACA | CCTATACCAG | CTCTGGACAA | 6120 |
| 5 | TTTAATTACC | TGACCTGGAT | TAGAACTGGA | AGCCCCAAGT | GCTCTCCTTC | AGACCTAGAT | 6180 |
| | TACCTAAAAA | TAAGTTTCAC | TGAGAAAAGGA | AAACAAGAAA | ATATCCTAAA | ATGGGTAAAT | 6240 |
| | GGTATGTCTT | GGGGAATGGT | ATATTATGGA | GGCTCGGGTA | AACAACCAGG | CTCCATTCTA | 6300 |
| 10 | ACTATTGCGC | TCAAAATAAA | CCAGCTGGAG | CCTCCAATGG | CTATAGGACC | AAATACGGTC | 6360 |
| | TTGACGGGTC | AAAGACCCCC | AACCCAAGGA | CCAGGACCAT | CCTCTAACAT | AACTTCTGGA | 6420 |
| 15 | TCAGACCCCA | CTGAGTCTAA | CAGCACGACT | AAAATGGGGG | CAAAACTTTT | TAGCCTCATC | 6480 |
| | CAGGGAGCTT | TTCAAGCTCT | TAAGTCCACG | ACTCCAGAGG | CTACCTCTTC | TTGTTGGCTA | 6540 |
| | TGCTTAGCTT | CGGGCCCAAC | TTACTATGAA | GGAATGGCTA | GAAGAGGGAA | ATTCAATGTG | 6600 |
| 20 | ACAAAAGAAC | ATAGAGACCA | ATGCACATGG | GGATCCCCAA | ATAAGCTTAC | CCTTACTGAG | 6660 |
| | GTTTCTGGAA | AAGGCACCTG | CATAGGAAAG | GTTCCCCCAT | CCCACCAACA | CCTTTGTAAC | 6720 |
| 25 | CACACTGAAG | CCTTTAATCA | AACCTCTGAG | AGTCAATATC | TGGTACCTGG | TTATGACAGG | 6780 |
| | TGGTGGGCAT | GTAATACTGG | ATTAACCCCT | TGTGTTTCCA | CCTTGCTTTT | TAACCAAAC | 6840 |
| | AAAGATTTT | GCATTATGGT | CCAAATTGTT | CCCCGAGTGT | ATTACTATCC | CGAAAAGCA | 6900 |
| 30 | ATCCTTGATG | AATATGACTA | CAGAAATCAT | CGACAAAAGA | GAGAACCCAT | ATCTCTGACA | 6960 |
| | CTTGCTGTGA | TGCTCGGACT | TGGAGTGGCA | GCAGTGCTAG | GAACAGGAAC | AGCTGCCCTG | 7020 |
| 35 | GTCACGGGAC | CACAGCAGCT | AGAAACAGGA | CTTAGTAACC | TACATCGAAT | TGTAACAGAA | 7080 |
| | GATCTCCAAG | CCCTAGAAAA | ATCTGTCAGT | AACCTGGAGG | AATCCCTAAC | CTCCTTATCT | 7140 |
| | GAAGTAGTCC | TACAGAATAG | AAGAGGGTTA | GATTTATTAT | TTCTAAAAGA | AGGAGGATTA | 7200 |
| 40 | TGTGTAGCCT | TGAAGGAGGA | ATGCTGTTTT | TATGTGGATC | ATTCAGGGGC | CATCAGAGAC | 7260 |
| | TCCATGAACA | AGCTTAGAGA | AAGGTTGGAG | AAGCGTCGAA | GGGAAAAGGA | AACTACTCAA | 7320 |
| 45 | GGGTGGTTTG | AGGGATGGTT | CAACAGGTCT | CTTTGGTTGG | CTACCCTACT | TTCTGCTTTA | 7380 |
| | ACAGGACCCT | TAATAGTCCT | CCTCCTGTTA | CTCACAGTTG | GGCCATGTAT | TATTAACAAG | 7440 |
| | TTAATTGCCT | TCATTAGAGA | ACGAATAAGT | GCAGTCCAGA | TCATGGTACT | TAGACAACAG | 7500 |
| 50 | TACCAAAGCC | CGTCTAGCAG | GGAAGCTGGC | CGCTAGCTCT | ACCAGTTCTA | AGATTAGAAC | 7560 |
| | TATTAACAAG | AGAAGAAGTG | GGGAATGAAA | GGATGAAAAT | ACAACCTAAG | CTAATGAGAA | 7620 |
| 55 | GCTTAAATT | GTTCTGAATT | CCAGAGTTTG | TTCCTTATAG | GTAAAAGATT | AGGTTTTTTG | 7680 |
| | CTGTTTTTAA | ATATGCGGAA | GTAAAATAGG | CCCTGAGTAC | ATGTCTCTAG | GCATGAAACT | 7740 |

5 TCTTGAAACT ATTTGAGATA ACAAGAAAAG GGAGTTTCTA ACTGCTTGTT TAGCTTCTGT 7800
AAAAC TGTT GCGCCATAAA GATGTTGAAA TGTGATACA CATATCTTGG TGACAACATG 7860
TCTCCCCCAC CCCGAAACAT GCGCAAATGT GTAAC TCTAA AACAATTTAA ATTAATTGGT 7920
CCACGAAGCG CGGGCTCTCG AAGTTTTTAA TTGACTGGTT TGTGATATTT TGAAATGATT 7980
10 GGT TTGTAAA GCGCGGGCTT TGT TGTGAAC CCCATAAAAG CTGTCCCGAC TCCACACTCG 8040
GGGCCGAGT CCTCTACCCC TGC GTGGTGT ACGACTGTGG GCCCCAGCGC GCTTGGGAATA 8100
AAAATCCTCT TGCTGTTTGC ATCAAAAAAA AA 8132

(2) INFORMATION FOR SEQ ID NO:4:

20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

30 TGCCTAGAGA CATGTACTC 19

(2) INFORMATION FOR SEQ ID NO:5:

35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

45 CCTCTTCTAG CCATTCCTTC A 21

(2) INFORMATION FOR SEQ ID NO:6:

50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

55 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TCGAGACTCG GTGGAAGGGC CC

22

(2) INFORMATION FOR SEQ ID NO:7:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

15 GGGCCCTTCC ACCGAGTCTC GA

22

(2) INFORMATION FOR SEQ ID NO:8:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

30 ACCTGGATCC ATGCATCCCA CG

22

(2) INFORMATION FOR SEQ ID NO:9:

- 35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CGTGGGATGC ATGGATCCAG GT

22

45 (2) INFORMATION FOR SEQ ID NO:10:

- 50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGCGCCACCT CCCGATTCCG

20

(2) INFORMATION FOR SEQ ID NO:11:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 10 (ii) MOLECULE TYPE: cDNA

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

15 CCGAATCGGG AGGTGGCGCC

20

(2) INFORMATION FOR SEQ ID NO:12:

- 20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 25 (ii) MOLECULE TYPE: cDNA

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

TCCCCTTAAG CTTGCCTCC

20

30 (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
35 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA

- 40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GGAGGCGAAG CTTAAGGGGA

20

45 (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
50 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA

- 55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

AAAAGCACAA AGGGCAGGAG AGC

23

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GCTCTCCTGC CCTTTGTGCT TTT

23

15 (2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CCTTTAGGAA CTGGTGGCC

20

30 (2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GGCCACCAGG TTCCTAAAGG

20

(2) INFORMATION FOR SEQ ID NO:18:

45

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

55

CCCCCAGATA TCCTCCATGC

20

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
5 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GCATGGAGGA TATCTGGGGG 20

(2) INFORMATION FOR SEQ ID NO:20:
15

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
20 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:
25
GCAGTTTCCA ATCAATCCCC AA 22

(2) INFORMATION FOR SEQ ID NO:21:
30

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
35 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:
40
TTGGGGATTG ATTGGAAACT GC 22

(2) INFORMATION FOR SEQ ID NO:22:
45

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:
50
TTTATGTTTG CCCAGGACCA CCA 23

55

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
5 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:
10 TGGTGGTCCT GGGCAAACAT AAA 23

(2) INFORMATION FOR SEQ ID NO:24:

15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
20 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:
25 GGGAGGTGGC GCCGGCTTAA CGT 23

(2) INFORMATION FOR SEQ ID NO:25:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:
40 ACGTTAAGCC GGCGCCACCT CCC 23

(2) INFORMATION FOR SEQ ID NO:26:

45 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:
55 CCCCCAACCC AAGGACCAGG ACCA 24

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

10 TGGTCCTGGT CCTTGGGTTG GGGG 24

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

GCAGCACGAC TAAAATGGGG GC 22

25 (2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29: —

GCCCCCATTT TAGTCGTGCT GC 22

40 (2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

50 CCCCCATCCC ACCAACACCT 20

(2) INFORMATION FOR SEQ ID NO:31:

55 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

AGGTGTTGGT GGGATGGGGG

20

10

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: cDNA

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

TCTCCCCAC CCCGAAACAT

20

25 (2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: cDNA

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

ATGTTTCGGG GTGGGGGAGA

20

40 (2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

50

AGCCAAGAAA GCCAGGTCCC CGAA

24

(2) INFORMATION FOR SEQ ID NO:35:

55 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

TTCGGGGACC TGGCTTTCTT GGCT

24

10 (2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

15 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

AGGCTCTGGT GCGGGTCTC C

21

25

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs

30 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

40 GGAGACCCGC CACCAGAGCC T

21

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

45 (A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

55

CCGCAGGGAT GGGTTTGGCA

20

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

15 TGCCAAACCC ATCCCTGCGG

20

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

30

GCTCACCTGG ACCCGACTGC CC

22

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

45

GGGCAGTCGG GTCCAGGTGA GC

22

(2) INFORMATION FOR SEQ ID NO:42:

50

(i) SEQUENCE CHARACTERISTICS:

- 55 (A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

5 GTTTACGGGA CGGGCAGCGA TGGC 24

(2) INFORMATION FOR SEQ ID NO:43:

10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: cDNA

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

GCCATCGCTG CCCGTCCCGT AAAC 24

(2) INFORMATION FOR SEQ ID NO:44:

25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: cDNA

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:44: _

TGGCTGGGGC GGCGGTGGTG GACGGG 26

40 (2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: cDNA

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

CCCGTCCACC ACCGCCGCC CAGCCA 26

55 (2) INFORMATION FOR SEQ ID NO:46:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- 10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:
GCCCCAAGCC CCAGAACCCA GACG 24
- 15 (2) INFORMATION FOR SEQ ID NO:47:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
20 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:
CGTCTGGGTT CTGGGGCTTT GGGC 24
- 30 (2) INFORMATION FOR SEQ ID NO:48:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
35 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- 40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:
45 GATGAACAGG CAGACATCTG 20
- (2) INFORMATION FOR SEQ ID NO:49:
(i) SEQUENCE CHARACTERISTICS:
50 (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 55 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

CGCTTACAGA CAAGCTGTGA

20

5

(2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

AGAACAAAGG CTGGGAAGC

19

20

(2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:

25

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

35

ATAGGAGACA GCCTGAACTC

20

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

40

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

50

GGACCATTGT CTGACCCTAT

20

(2) INFORMATION FOR SEQ ID NO:53:

55

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: cDNA

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

GTCAACACCT ATACCAGCTC

20

(2) INFORMATION FOR SEQ ID NO:54:

15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: cDNA

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

CATCTGAGGT ATAGCAGGTC

20

(2) INFORMATION FOR SEQ ID NO:55:

30

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: cDNA

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

GCAGGTGTAG GAACAGGAAC

20

45 (2) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: cDNA

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

ACCTGTTGAA CCATCCCTCA 20

(2) INFORMATION FOR SEQ ID NO:57:

5 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

CGAATGGAGA GATCCAGGTA 20

(2) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

25 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

CCTGCATCAC TTCTCTTACC 20

(2) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

40 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

50 TTGCCTGCTT GTGGAATACG 20

(2) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:

55 (A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

10 CAAGAGAAGA AGTGGGGAAT G

21

(2) INFORMATION FOR SEQ ID NO:61:

(i) SEQUENCE CHARACTERISTICS:

15

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

25 CACAGTCGTA CACCACGCAG

20

(2) INFORMATION FOR SEQ ID NO:62:

(i) SEQUENCE CHARACTERISTICS:

30

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

40

GGGAGACAGA AGAAGAAAGG

20

(2) INFORMATION FOR SEQ ID NO:63:

(i) SEQUENCE CHARACTERISTICS:

45

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: cDNA

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

CGATAGTCAT TAGTCCCAGG

20

(2) INFORMATION FOR SEQ ID NO:64:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 10 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

15 TGCTGGTTTG CATCAAGACC G 21

(2) INFORMATION FOR SEQ ID NO:65:

- 20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 25 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

30 GTCGCAAAGG CATACTGCT 20

(2) INFORMATION FOR SEQ ID NO:66:

- 35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
40 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

45 ACAGAGCCTC TGCTAAGAAG 20

(2) INFORMATION FOR SEQ ID NO:67:

- 50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
55 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

GCAGCTGTTG ACAATCATC

19

10 (2) INFORMATION FOR SEQ ID NO:68:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

TATGAGGAGA GGGCTTGACT

20

25 (2) INFORMATION FOR SEQ ID NO:69:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
30 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

AGCAGACGTG CTAGGAGGT

40

19

(2) INFORMATION FOR SEQ ID NO:70:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
45 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

55 TCCTCTTGCT GTTTGCATC

19

(2) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
5 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

15 CAGACACTCA GAACAGAGAC 20

(2) INFORMATION FOR SEQ ID NO:72:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
20 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

30 ACATCGTCTA ACCCACCTAG 20

(2) INFORMATION FOR SEQ ID NO:73:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
35 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

45 CTCGTTTCTG GTCATACCTG A 21

(2) INFORMATION FOR SEQ ID NO:74:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 base pairs
50 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

WO 97/21836

- 80 -

PCT/US96/19680

GAGTACATCT CTCTAGGCA

19

What is claimed is:

1. A purified nucleic acid which can specifically hybridize with the sequence of SEQ ID NO: 1 or its complement, provided that said nucleic acid is other than the entire retroviral genome of SEQ ID NO: 1 or its complement.
5
2. The purified nucleic acid of claim 1, wherein said nucleic acid is at least one nucleotide longer, or at least 1 nucleotide shorter, or differs in sequence at at least one position from SEQ ID NO: 1 or its complement.
10
3. The purified nucleic acid of claim 1 wherein said nucleic acid has at least 72% sequence identity or homology with a sequence from SEQ ID NO: 1 or its complement.
4. The purified nucleic acid of claim 1, wherein said nucleic acid is at least 15 nucleotides in length.
15
5. The purified nucleic acid of claim 1, wherein said nucleic acid can specifically hybridize with a translatable region of the retroviral genome of SEQ ID NO: 1, or its complement.
20
6. The purified nucleic acid of claim 1, wherein said nucleic acid can specifically hybridize with a region from the gag, pol, or env gene.
7. The purified nucleic acid of claim 1, wherein said nucleic acid can specifically hybridize with an untranslated region of the retroviral genome of SEQ ID NO: 1, or its complement.
25
8. The purified nucleic acid of claim 1, wherein said nucleic acid can specifically hybridize with a non-conserved region of the retroviral genome of SEQ ID NO: 1, or its complement.
30
9. The purified nucleic acid of claim 1, wherein said nucleic acid can specifically hybridize with highly conserved regions of the retroviral genome of SEQ ID NO: 1, or its complement.
35
10. The purified nucleic acid of claim 1, wherein the nucleic acid is selected from the group consisting of SEQ ID NOs: 4-74.

11. A purified nucleic acid which hybridizes under stringent conditions to a nucleic acid chosen from: a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 of SEQ ID NO:1, or naturally occurring mutants thereof, a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from
5 nucleotides 4871-8060 of SEQ ID NO:1, or naturally occurring mutants thereof, and a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 of SEQ ID NO:1, or naturally occurring mutants thereof.

12. The purified nucleic acid of claim 11, wherein said nucleic acid is a nucleic
10 acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 of SEQ ID NO:1, or naturally occurring mutants thereof.

13. The purified nucleic acid of claim 11, wherein said nucleic acid is a nucleic
15 acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, or naturally occurring mutants thereof.

14. The purified nucleic acid of claim 11, wherein said nucleic acid is a nucleic
20 acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 of SEQ ID NO:1, or naturally occurring mutants thereof.

15. A reaction mixture which includes a target nucleic acid and a second nucleic acid, wherein the second nucleic acid is chosen from: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically
25 hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g, from nucleotides 3112-4683) of SEQ ID NO:1,
30 nucleotides 598-2169 (e.g, from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof;

a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or
35 antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof;

a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g. from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g. from nucleotides 4738-6722) of SEQ ID NO:2, or
5 nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; a swine or miniature swine retroviral nucleic acid; or a Tsukuba nucleic acid.

10 16. A purified nucleic acid which can specifically hybridize with the sequence of SEQ ID NO: 2 or its complement.

17. The purified nucleic acid of claim 16, wherein said nucleic acid is at least one nucleotide longer, or at least 1 nucleotide shorter, or differs in sequence at at least one position from SEQ ID NO: 2 or its complement.

15 18. The purified nucleic acid of claim 16, wherein said nucleic acid has at least 72% sequence identity or homology with a sequence from SEQ ID NO: 2 or its complement.

20 19. The purified nucleic acid of claim 16, wherein said nucleic acid is at least 15 nucleotides in length.

20. The purified nucleic acid of claim 16, wherein said nucleic acid can specifically hybridize with a region from the gag, pol. or env gene.

25 21. The purified nucleic acid of claim 16, wherein said nucleic acid hybridizes under stringent conditions to a nucleic acid chosen from: a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 598-2169 of SEQ ID NO:2, or naturally occurring mutants thereof, a nucleic acid of at least 10 consecutive
30 nucleotides of sense or antisense sequence from nucleotides 2320-4737 of SEQ ID NO:2, or naturally occurring mutants thereof, and a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4738-6722 of SEQ ID NO:2, or naturally occurring mutants thereof.

35 22. A purified nucleic acid which can specifically hybridize with the sequence of SEQ ID NO: 3 or its complement.

23. The purified nucleic acid of claim 22, wherein said nucleic acid is at least one nucleotide longer, or at least 1 nucleotide shorter, or differs in sequence at at least one position from SEQ ID NO: 3 or its complement.

5 24. The purified nucleic acid of claim 22, wherein said nucleic acid has at least 72% sequence identity or homology with a sequence from SEQ ID NO: 3 or its complement.

10 25. The purified nucleic acid of claim 22, wherein said nucleic acid is at least 15 nucleotides in length.

26. The purified nucleic acid of claim 22, wherein said nucleic acid can specifically hybridize with a region from the gag, pol, or env gene.

15 27. The purified nucleic acid of claim 22, wherein said nucleic acid hybridizes under stringent conditions to a nucleic acid chosen from: a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof, a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 585-2156 of SEQ ID NO:3, or
20 naturally occurring mutants thereof, and a nucleic acid of at least 3 consecutive nucleotides of sense or antisense sequence from nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof.

25 28. A method for screening a cell or a tissue for the presence or expression of a swine or miniature swine retrovirus comprising:

 contacting a target nucleic acid from the tissue with a second nucleic acid selected from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID
30 NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g, from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g, from
35 nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence

from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense
5 sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof, under conditions in which hybridization can occur, hybridization being indicative of the presence or expression of an endogenous swinw or miniature swine retrovirus or retroviral sequence in the tissue.

10

29. A method for screening a swine or miniature swine genome for the presence of a porcine retrovirus, comprising:

contacting the miniature swine genomic DNA with a second nucleic acid selected from the group of: a sequence which can specifically hybridize to a porcine retroviral
15 sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10
20 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g, from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g, from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof;

a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which
25 encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof;

a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which
30 encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof, under conditions in which the sequences can hybridize,

35 hybridization being indicative of the presence of the endogenous porcine retroviral sequence in the miniature swine genome.

30. A method of assessing the potential risk associated with the transplantation of a graft from a donor swine or miniature swine into a recipient animal, comprising:

- contacting a target nucleic acid from the donor, recipient or the graft, with a second nucleic acid selected from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g, from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g, from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof, under conditions in which the sequences can hybridize, hybridization being indicative of a risk associated with the transplantation.

31. A method of providing a swine or miniature swine free of an activatable retrovirus insertion at a preselected site, comprising:

- performing a cross between a first miniature swine having a retroviral insertion at the preselected site and a second miniature swine not having a retroviral insertion at a preselected site, and recovering a progeny miniature swine, not having the insertion, wherein the presence or absence of the retroviral insertion is determined by contacting the genome of a miniature swine with a nucleic acid chosen from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense

- sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g, from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g, from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or
- 5 naturally occurring mutants thereof;
- a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring
- 10 mutants thereof;
- a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or
- 15 nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof.

32. A method of localizing the origin of a porcine retroviral infection, comprising:
- contacting a target nucleic acid from the graft or organ with a second nucleic acid selected from the group of: a sequence which can specifically hybridize to a porcine
- 20 retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of
- 25 at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g, from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g, from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g, from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof;
- a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which
- 30 encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof;
- a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which
- 35 encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof;

contacting a target nucleic acid from the recipient with a second nucleic acid selected from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g. from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g. from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g. from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof;

a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof;

a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g. from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g. from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof; hybridization to the nucleic acid from the graft correlates with the porcine retroviral infection in the graft; and hybridization to the nucleic acid from the recipient correlates with the porcine retroviral infection in the recipient.

25

33. A method of screening a human subject for the presence or expression of an endogenous porcine retrovirus comprising:

contacting a target nucleic acid derived from the human subject with a second nucleic acid selected from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g. from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g. from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g. from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof;

- a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737 of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof;
- a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g. from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g. from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof, under conditions in which the sequences can hybridize, hybridization being indicative of the presence of the endogenous porcine retrovirus or retroviral sequences in the human subject.
34. A transgenic miniature swine having a transgenic element at an endogenous porcine retroviral insertion site which corresponds to the retroviral genome of SEQ ID NO: 1, 2, or 3, and wherein said element alters the activity of the endogenous porcine retrovirus.
35. A method of detecting a recombinant virus or other pathogen, comprising:
providing a pathogen having porcine retroviral sequence; and
determining if the pathogen includes non-porcine retroviral sequence, the presence of non-porcine retroviral sequence being indicative of viral recombination.
36. A method of determining the copy number, size, or completeness of a porcine retrovirus, comprising:
contacting a target nucleic acid from the donor, recipient or a graft, with a second nucleic acid selected from the group of: a sequence which can specifically hybridize to a porcine retroviral sequence; a sequence which can specifically hybridize to the sequence of SEQ ID NO:1 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:2 or its complement; a sequence which can specifically hybridize to the sequence of SEQ ID NO:3 or its complement; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a gag protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2452-4839 (e.g. from nucleotides 3112-4683) of SEQ ID NO:1, nucleotides 598-2169 (e.g. from nucleotides 598-2169) of SEQ ID NO:2, or nucleotides 585-2156 (e.g. from nucleotides 585-2156) of SEQ ID NO:3, or naturally occurring mutants thereof;
a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a pol protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 4871-8060 of SEQ ID NO:1, nucleotides 2320-4737

of SEQ ID NO:2, or nucleotides 2307-5741 of SEQ ID NO:3, or naturally occurring mutants thereof;

- 5 a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence which encodes a env protein; a nucleic acid of at least 10 consecutive nucleotides of sense or antisense sequence from nucleotides 2-1999 (e.g, from nucleotides 86-1999) of SEQ ID NO:1, nucleotides 4738-6722 (e.g, from nucleotides 4738-6722) of SEQ ID NO:2, or nucleotides 5620-7533 of SEQ ID NO:3, or naturally occurring mutants thereof.

- 10 37. A method for screening a tissue for the presence or expression of a swine or a miniature swine retroviral sequence comprising:

contacting a tissue sample with an antibody specific for a retroviral protein, thereby determining if the sequence is present or expressed.

- 15 38. A purified nucleic acid which can specifically hybridize to a nucleic acid sequence comprising nucleotides 2-1999 of SEQ ID NO:1, nucleotides 4871-8060 of SEQ ID NO:1, or nucleotides 2452-4839 of SEQ ID NO:1.

CTUGAGACTC GGTGAAGGG CCCATTATCTC GTACTTTTGA CCACACCAAC 50 (SEQ ID NO: 1)
GGCTGTGAAA GTCGAAGGAA TCTCCACCTG GATCCATGCA TOCCACGTTA 100
AGCCGGCGCC ACCTCCCGAT TCGGGGTGGA AAGCCGAAAA GACTGAAAAT 150
CCCTTAAGC TTGCGCTCCA TCGGTGGTT CTTACTCTG TCAATAACCT 200
CTCAGACTAA TGGTATGCC ATAGGAGACA GCGTGAATC CCATAAACCC 250
TTATCTCTCA CTTGGTTAAT TACTGACTCC GGCACAGGTA TTAATATCAA 300
CAACACTCAA GGGGAGGCTC CTTTAGGAAC CTGGTGGCCT GATCTATAAG 350
TTTGCCCTCAG ATCAGTTATT CTTAGTCTGA CCTCACCCCC AGATATCCTC 400
CATGCTCAGG GATTTTATGT TTGCCAGGA CCACCAAATA ATGGAACA 450
TTGCGGAAT CCCAGAGATT TCTTTTGTA ACAATGGAAC TGTGTAACCT 500
CTAATGATGG ATATTGAAA TGCCCAACCT CTCAGCAGGA TAGGGTAAGT 550
TTTTCTTATG TCAACACCTA TACCAGCTCT GGACAATTTA ATTACCTGAC 600
CTGGATTAGA ACTGGAAGCC CCAAGTGCTC TCCTTCAGAC CTAGATTACC 650
TAAAAATAAG TTTCACCTGAG AAAGGAAAAC AAGAAATAT CCTAAAATGG 700
GTAAATGGTA TGTCTTGGG AATGGTATAT TATGGAGGCT CGGGTAAACA 750
ACCAGGCTCC ATTCTAACTA TTGCGCTCAA AATAAACCCAG CTGGAGCCTC 800
CAATGGCTAT AGGACCAAAT ACGGTCTTGA CGGGTCAAAG ACCCCCAACC 850
CAAGGACCAG GACCATCCTC TAACATAACT TCTGGATCAG ACCCCACTGA 900
GTCTAGCAGC ACGACTAAAA TGGGGGCAA ACTTTTITAGC CTCATCCAGG 950
GAGCTTTTCA AGCTCTTAAC TCCACGACTC CAGAGGCTAC CTCTCTTGT 1000
TGGCTATGCT TAGCTTTGGG CCCACCTTAC TATGAAGGAA TGGCTAGAAG 1050
AGGGAAATTC AATGTGACAA AAGAACATAG AGACCAATGC ACATGGGGAT 1100
CCCAAATAA GCTTACCCCT ACTGAGGTTT CTGGAAGG CACCTGCATA 1150
GGAAAGGTTT CCCCATCCCA CCAACACCTT TGTAACCACA CTGAAGCCCT 1200
TAATCAAACC TCTGAAAGTC AATATCTGGT ACCTGGTTAT GACAGGTGGT 1250
GGGCATGTA TACTGGATTA ACCCCTTGTG TTTCACCTT GGTTTTAAAC 1300

FIGURE 1

CAAACATAAG ATTTTTGCAAT TATGGTCCAA ATTGTTCCCC GAGTGTATTA 1350 (SEQ ID NO: 1)
cont'd
CTATCCCGAA AAAGCAATCC TTGATGAATA TGAATACAGA AATCATCGAC 1400
AAAAGAGAGA ACCCATATCT CTGACACTTG CTGTGATGCT CGGACTTGGG 1450
GTGGCAGCAG GTGTAGGAAC AGGAACAGCT GCCCTGGTCA CCGGACCACA 1500
GCAGCTAGAA ACAGGACTTA GTAACCTACA TCGAATTGTA ACAGAAGATC 1550
TCCAGCCCT AGAAAAATCT GTCAGTAACC TGGAGGAATC CCTAACCTCC 1600
TTATCTGAAG TAGTCTACA GAATAGAAGA GGTTAGATT TATTATTCT 1650
AAAAGAAGGA GGATTATGTG TAGCCTTGAA GGAGGAATGC TGTTTTTATG 1700
TGGATCATTC AGGGCCATC ACAGACTCCA TGAACAACT TAGAGAAAGG 1750
TTGGAGAAGC GTCGAAGGA AAAGGAACT ACTCAAGGT GGTTTGAGG 1800
ATGTTTCAAC AGGTCTCTT GGTGGCTAC CCTACTTCT GCTTTAACAG 1850
GACCTTAAT AGTCTCTC CTGTTACTCA CAGTTGGGC ATGTATTATT 1900
AACAAGTTAA TTGCTTCAT TAGAGAACGA ATAAGTGCAG TCCAGATCAT 1950
GGTACTTAGA CAACAGTACC AAAGCCGTC TAGCAGGAA GCTGGCCCT 2000
AGCTCTACCA GTTCTAAGAT TAGAACTATT AACAAGAGAA GAAGTGGGA 2050
ATGAAAGGAT GAAAATACAA CCTAAGCTAA TGAGAAGCTT AAAATTGTTC 2100
TGAATTCCAG AGTTTGTTC TTATAAGTAA AAGATTAGGT TTTTTCCT 2150
TTTAAATAT GCGGAAGTAA AATAGCCCT GAGTACATGT CTCTAGGCAT 2200
GAACTTCTT GAACTATTT GAGATAACAA GAAAAGGAG TTTCTAACTG 2250
CTTGTTAGC TTCTGTAAAA CTGGTTGCC CATAAAGATG TTGAAATGTT 2300
GATACACATA TCTTGGTGAC AACATGTCT CCCCCCCC AAACATGCC 2350
AAATGTGTAA CTCTAAAACA ATTTAAATTA ATTGGTCCAC GAAGCGCGG 2400
CTCTCGAAGT TTTAAATTGA CTGGTTGTG ATATTTTGAA ATGATTGGTT 2450
TGTAAGGCG GGGCTTTGCT GTGAACCCCA TAAAGCTGT CCGACTCCA 2500
CACTCGGGC CGCAGTCTC TACCCCTGCG TGGTGTACGA CTGTGGG 2550

FIGURE 1, CONT.

3/34

| | | | | | | |
|------------|-------------|------------|-------------|------------|------|----------------|
| CAGCGCGCTT | GGAAATAAAAA | TCCTCTTGCT | GTTTGCAATCA | AGACCGCTTC | 2600 | (SEQ ID NO: 1) |
| TCGTGAGTGA | TTAAGGGGAG | TCGCCTTTTC | CGAGCCTGGA | GGTCTTTTTT | 2650 | cont'd |
| GCTGGTCTTA | CATTTGGGGG | CTCGTCCGGG | ATCTGTGGCG | GCCACCCCTA | 2700 | |
| ACACCCGAGA | ACCGACTTGG | AGGTAAAAAG | GATCCTCTTT | TTAACGTGTA | 2750 | |
| TGCAATGACC | GGCCGGGGTC | TCTGTCTGTA | GTGTCTGTTT | TCAGTGGTGC | 2800 | |
| CGCCTTTTCG | TTTGCACTG | TCCTCTCAGG | CGTAAGGGC | TGGGGGACTG | 2850 | |
| TGATCAGCAG | ACGTGCTAGG | AGGATCACAG | GCTGCTGCCC | TGGGGGACGC | 2900 | |
| CCCGGAGGT | GAGGAGAGCC | AGGGACCGCT | GGTGGTCTCC | TACTGTCCGT | 2950 | |
| CAGAGGACCG | AATTCTGTTG | CTGAAGCGAA | AGCTTCCCCC | TCCGCGACCG | 3000 | |
| TCCGACTCTT | TTGCCTGCTT | GTGGAATACG | TGGACGGGTC | ACGTGTGICT | 3050 | |
| GGATCTGTTG | GTTCCTGTTT | TGTGTGCTTT | TGTCTGTGTT | GTCTTGTGCT | 3100 | |
| ACAGTTTAA | TATGGGACAG | ACGGTGACGA | CCCTCTTAG | TTTCACTCTC | 3150 | |
| GACCATTCGA | CTGAAGTTAA | ATCCAGGGCT | CATAATTTGT | CAGTTCAGGT | 3200 | |
| TAAGAAGGGA | CCTTGGCAGA | CTTCTGTGTT | CTCTGAATGG | CCGACATTCC | 3250 | |
| ATGTTGGATG | GCCATCAGAG | GGACCTTTA | ATTCTGAGAT | TATCCTGGCT | 3300 | |
| GTTAAAGCAA | TTATTTTICA | GACTGGACCC | GGCTCTCATT | CCGATCAGGA | 3350 | |
| GCCCTATATC | CTTACGTGGC | AAGATTTGGC | AGAGGATCCT | CCGCCATGGG | 3400 | |
| TTAAACCATG | GCTGAATAAG | CCAAGAAAGC | CAGCTCCCCG | AATTCTGGCT | 3450 | |
| CTTGGAGAGA | AAAACAAACA | CTCGGCTGAA | AAAGTCAAGC | CCTCTCCTCA | 3500 | |
| TATCTACCCC | GAGATTGAGG | AACCACCGGC | TTGGCCGGAA | CCCCAATCTG | 3550 | |
| TTCCCCCACC | CCCTTATCTG | GCACAGGGTG | CCGCGAGGGG | ACCTTTTGCC | 3600 | |
| CCTCCTGGAG | CTCCGGCGGT | GGAGGGACCT | TCTGCAGGGA | CTCGGAGCCG | 3650 | |
| GAGGGGCGCC | ACCCCGGAGC | GGACAGACGA | GATCGCGACA | TTACCGCTGC | 3700 | |
| GCACCTACCG | CCCTCCACAA | CCGGGGGGCC | AAATTCAGCC | CCTCCAGTAT | 3750 | |
| TGGCCCTTTT | CTTCTGCAGA | TCTCTATAAT | TGGAAACTA | ACCATCCCCC | 3800 | |

FIGURE 1, CONT.

TTTCTCGGAG GATCCCCAAC GCTTCACGGG GTTGGTGGAG TCCCTTATGT 3850 (SEQ ID NO: 1)
TCTCTACCA GCTTACTTGG GATGATTGTC AACAGCTGCT CCAGACATC cont'd
TTCACAACCG AGGAGCGAGA GAGAATTCTA TTAGAGGCTA GAAAAAATGT 3900
TCCCTGGGGCC GACGGGGGAC CCACGGGTTT GCAAAATGAG ATTGACATGG 3950
GATTTCCCTT AACTCGCCCC GGTITGGGACT ACAACACGGC TGAAGGTAGG 4000
GAGAGCTTGA AAATCTATCG CCAGGCTCTG GTGGCGGGTC TCCGGGGCGC 4050
CTCAAGACGG CCCACTAATT TGGCTAAGGT AAGAGAAGTG ATGCAGGGAC 4100
CGAATGAACC CCCCTCTGTT TTTCTTGAGA GGCTCTTGGA AGCCTTCAGG 4150
CGGTACACCC CTTTTGATCC CACCTCAGAG GCCCCAAAAG CCTCAGTGGC 4200
TTTGGCCTTT ATAGGACAGT CAGCCTTGGA TATTAGAAAG AAGCTTCAGA 4250
GACTOGAAGG GTTACAGGAG GCTGAGTTAC GTGATCTAGT GAAAGAGGCA 4300
GAGAAAGTAT ATTACAAAAG GGAGACAGAA GAAGAAAGGG AACAAAGPAA 4350
AGAGAGAGAA AGAGAGGAAA GGGAGGAAAG ACGTAATAAA CCGCAAGAGA 4400
AGAATTTGAC TAAGATCTTG GCTGCAGTGG TTGAAGGGAA AAGCAATACG 4450
GAAAGAGAGA GAGATTTTAG GAAACTTAGG TCAGGCCCTA GACAGTCAGG 4500
GAACCTGGGC AATAGGACCC CACTCGACAA GGACCAATGT GCATATTGTA 4550
AAGAAAGAGG ACACTGGGCA AGGAACTGCC CCAAGAAGGG AAACAAAGGA 4600
CCAAGGATCC TAGCTCTAGA AGAAGATAAA GATTAGGGGA GACGGGGTTC 4650
GGACCCCCCTC CCCGAGCCCA GGGTAACTTT GAAGGTGGAG GGGCAACCAG 4700
TTGAGTTTCCT GGTTGATACC GGAGCGAAAC ATTCACTGCT ACTACAGCCA 4750
TTAGGAAAAC TAAAAGATAA AAATCCTGG GTGATGGGTG CACAGGGCAA 4800
CAACAGTATC CATGGACTAC CCGAAGACAG TTGACTTGGG AGTGGGACGG 4850
GTAACCCACT CGTTTCTGGT CATACCTGAG TCCCCAGCAC CCCTCTTAGG 4900
TAGAGACTTA TTGACCAAGA TGGGAGCACA AATTTCTTTT GAACAAGGA 4950
AACCAGAAGT GTCTGCAAAT AACAAACCTA TCACTGTGTT GACCCCTCAA 5000

FIGURE 1, CONT.

5/34

| | | |
|--|------|----------------|
| TTAGATGACG AATATCGACT ATACTCTCCC CTAGTAAAGC CTGATCAAAA | 5100 | (SEQ ID NO: 1) |
| TATACAATTC TGGTTGGAAC AGTTTCCCCA AGCCTGGGCA GAAACCGCAG | 5150 | cont'd |
| GGATGGGTTT GCCAAAGCAA GTTCCCCCAC AAGTTATTC AACTGAAGGCC | 5200 | |
| AGTCCCACAC CAGTGTGAGT CAGACAGTAC CCCTTGAGTA AAGAAGCTCA | 5250 | |
| AGAAGGAATT CGGCCCATG TCCAAAGATT AATCCAACAG GGCATCCTAG | 5300 | |
| TTCTGTCCA ATCTCCCTGG AATACTCCCC TGCTACCGGT TAGAAAGCCT | 5350 | |
| GGGACTAATG ACTATCGACC AGTACAGGAC TTGAGAGAGG TCAATAAAGC | 5400 | |
| GGTGCAGGAT ATACACCCAA CAGTCCCGAA CCCTTATAAC CTCTTGTTG | 5450 | |
| CTCTCCACC CCAACGGAGC TGGTATACAG TATTGGACTT AAAGGATGCC | 5500 | |
| TTCTTCGGC TGAGATTACA CCCACTAGC CAACCACTTT TTGCTTGA | 5550 | |
| ATGGAGAGAT CCAGGTACCG GAAGAACCG GCAGCTCACC TGGACCCGAC | 5600 | |
| TGCCCCAAGG GTTCAAGAAC TCCCCGACCA TCTTTGACGA AGCCCTACAC | 5650 | |
| AGAGACCTGG CCAACTTCAG GATCCAACAC CCTCAGGTGA CCTCTCTCA | 5700 | |
| GTACGTGGAT GACCTGCTTC TGGGGGAGC CACCAACAG GACTGCTTAG | 5750 | |
| AAGGCACGAA GGCCTACTG CTGGAATTGT CTGACCTAGG CTACAGAGCC | 5800 | |
| TCTGCTAAGA AGGCCAGAT TTGCAGGAGA GAGGTACAT ACITGGGGTA | 5850 | |
| CAGTTTACGG GACGGGCAGC GATGGCTGAC GGAGGCACCG AAGAAAACG | 5900 | |
| TAGTCCAGAT ACCGGCCCCA ACCACAGCCA AACAAATGAG AGAGTTTTTG | 5950 | |
| CGGACAGCTG GATTTTGCAG ACTGTGGATC CCGGGGTTTG CGACCTTAGC | 6000 | |
| AGCCCCACTC TACCCGCTAA CCAAAGAAAA AGGGGAATTC TCCTGGGCTC | 6050 | |
| CTGAGCACCA GAAGGCATTT GATGCTATCA AAAAGGCGCT GCTGAGCGCA | 6100 | |
| CCTGCTCTGG CCTCCTTGA CGTAACTAAA CCTTTTACCC TTTATGTGGA | 6150 | |
| TGAGCGTAAG GGAGTAGCCC GGGGAGTTTT AACCCAAACC CTAGGACCAT | 6200 | |
| GGAGAAGACC TGTGCGCTAC CTGTCAAAGA AGCTCGATCC TGTAGCCAGT | 6250 | |
| GGTTGGCCCA TATGCTGAA GGCTATCGCA GCTGTGGCCA TACTGTCAA | 6300 | |

FIGURE 1. CONT.

| | | |
|--|------|----------------|
| GGACGCTGAC AAATTGACTT TGGACAAGA ATATAACTGT AATAGCCCCC | 6350 | (SEQ ID NO: 1) |
| CATGCATTGG AGAACATCGT TGGCAGCCC CCAGACCGAT GGATGACCAA | 6400 | cont'd |
| CGCCCGCATG ACCCACTATC AAAGCCTGCT TCTCACAGAG AGGGTCACGT | 6450 | |
| TGGCTCCACC AACCGCTCTC AACCTGCCA CTCTTCTGCC TGAAGAGACT | 6500 | |
| GATGAACCAG TGAATCATGA TTGCCATCAA CTATTGATTG AGGAGACTGG | 6550 | |
| GGTCCGCAAG GACCTTACAG ACATACCGCT GACTGGAGAA GTGCTAACCT | 6600 | |
| GGTTCACTGA CCGAAGCAGC TATGTGGTGG AAGGTAAGAG GATGGCTGGG | 6650 | |
| GCGGGGGTGG TGGACGGGAC CCGCAGGATC TGGCCAGCA GCCTGCCGGG | 6700 | |
| AGGAACCTCA GCACAAAAGG CTGAGCTCAT GGCCCTCAGC CAAGCTTTGC | 6750 | |
| GGCTGGCCGA AGGGAAATCC ATAAACATTT ATACGGACAG CAGGTATGCC | 6800 | |
| TTTGGGACTG CACACGTACA TGGGGCCATC TATAAACAAA GGGGGTTGCT | 6850 | |
| TACCTCAGCA GGGAGGGAAA TAAAGAACA AGAGGAAATT CTAAGCCTAT | 6900 | |
| TAGAAGCCGT ACATTTACCA AAAAGGCTAG CTATTATACA CTGTCTGGA | 6950 | |
| CATCAGAAAG CTAAAGATCT CATATCCAGA GGAAACCAGA TGGCTGACCG | 7000 | |
| GGTTGCCAAG CAGGCAGCCC AGGGTGTTAA CCTTCTGCTT ATAATAGAAA | 7050 | |
| TGCCCCAAGC CCCAGAACCC AGACGACAGT ACACCTPAGA AGACTGGCAA | 7100 | |
| GAGATAAAAA AGATAGACCA TTCTCTGAGA CTCCGGAAGG GACCTGCTAT | 7150 | |
| ACCTCAGATG GGAAGGAAAT CCTGCCCCAC AAAGAAGGGT TAGAATATGT | 7200 | |
| CCAACAAGAT ACATCGTCTA ACCACCTAG GAACTAAACA CCTGCAGCAG | 7250 | |
| TTGGTCAGAA CATCCCCCTA TCATGTCTTG AGGCTACCAG GAGTGGCTGA | 7300 | |
| CTCGGTGGIC AAACATTGTG TGCCCTGCCA GCTGGTTAAT GCTAATCCTT | 7350 | |
| CCAGAATGCC TCCAGGGAAG AGACTAAGGG GAAGCCACCC AGGCGCTCAC | 7400 | |
| TOGGAAGTGG ACTTCACTGA GGTAAAGCCG GCTAAATATG GAAACAAATA | 7450 | |
| CCTATTGGTT TTTGTAGACA CCTTTTCAGG ATGGGTAGAG GCTTATCCTA | 7500 | |
| CTAAGAAAGA GACTTCAACC GTGGTAGCTA AAAAAATACT GGAAGAAATT | 7550 | |

FIGURE 1, CONT.

7/34

| | | |
|--|------|----------------|
| TTTCCAAGAT TTGGAATACC TAAAGTAATA OGGTCAGACA ATGGTCCAGC | 7600 | (SEQ ID NO: 1) |
| TTTTGTGTGCC CAGGTAAGTC AOGGACTGOC CAAGATATTTG GGGATTGATT | 7650 | cont'd |
| GGAAACTGCA TTGTGCATAC AGACCCCAAA GCTCAGGACA GGTAGAGAGG | 7700 | |
| ATGAATAGAA CCATTAAAGA GACCCCTTACT AAATTGACCG CGGAGACTGG | 7750 | |
| CGTTAATGAT TGGATAGCTC TCCTGCCCTT TGTGCTTTTT AGGGTTAGGA | 7800 | |
| ACACCCCTCG ACAGTTTGGG CTGACCCCTT ATGAATTACT CTACGGGGGA | 7850 | |
| CCCCCCCCAT TGGTAGAAAT TGCTTCTGTA CATAGTGCTG ATGTGCTGCT | 7900 | |
| TTCCAGCCT TTGTTCTCTA GGCTCAAGCC ACTTGAGTGG GTGAGACAAC | 7950 | |
| GAGCGTGGAG GCAACTCCGG GAGGCCTACT CAAGAGGAGG AGACTTGCG | 8000 | |
| ATCCACATC GTTCCAAGT GGGAGATTCA GTCTACGTTA GACCCACCC | 8050 | |
| TGCAGGAAAC | 8060 | |

FIGURE 1, CONT.

8/34

(SEQ ID NO: 2)

```

      10      20      30      40      50      60
      *      *      *      *      *      *
CTACCCCTGC GTGGTGTACG ACTGTGGGCC CCAGCGCGCT TCGAATAAAA ATCCTCTTGC

      70      80      90     100     110     120
      *      *      *      *      *      *
TCTTTGCATC AAGACCGCTT CTGTGAGTGT ATTTGGGGTG TCCCTCTTTC CGAGCCCGGA

      130     140     150     160     170     180
      *      *      *      *      *      *
CGAGGGGGAT TGTTCCTTTA CTCGCCCTTC ATTTGGTGGG TTGGCCCGGA AATCCTGGGA

      190     200     210     220     230     240
      *      *      *      *      *      *
CCACCCCTTA CACCCGAGAA CCGACTTGA GGTAAAGGA TCCCTTTGG AACATATGTG

      250     260     270     280     290     300
      *      *      *      *      *      *
TGTGTGGGCC GGCGTCTCTG TCTGAGTGT CTGTTTGGG TGATCCGCC TTTCCGTTTG

      310     320     330     340     350     360
      *      *      *      *      *      *
CAGCTCTCCT CTCAGACCGT AAGGACTTGA GGACTGTGAT CAGCAGACGT GCTAGGAGGA

      370     380     390     400     410     420
      *      *      *      *      *      *
TCACAGGCTG CCACCTTGGG GGACGCCCGG GGAGGTGGGG AGAGCCAGGG ACGCCTGGTG

      430     440     450     460     470     480
      *      *      *      *      *      *
GTCTCTACT GTCCGTCAGA GGACCGAGTT CTGTTGTGA AGCGAAAGCT TCCCTCTCCG

      490     500     510     520     530     540
      *      *      *      *      *      *
CGGCGGTCCG ACTCTTTTGC CTGCTTGTGG AAGACGCGGA CCGGTGCGGT GTCTCTGGAT

      550     560     570     580     590     600
      *      *      *      *      *      *
CTGTGGGTTT CTGTTTGGTG TGTCTTTGTC TTGTGGGTCC TTGTCTACAG TTCTAAT ATG
                                         Met>

      610     620     630     640
      *      *      *      *
GGA CAG ACA GTG ACT ACC CCC CTT AGT TTG ACT CTC GAC CAT TGG ACT
Gly Gln Thr Val Thr Thr Pro Leu Ser Leu Thr Leu Asp His Trp Thr>

      650     660     670     680     690
      *      *      *      *      *
GAA GTT AGA TCC AGG CCT CAT AAT TTG TCA GTT CAG GTT AAG AAG GGA
Glu Val Arg Ser Arg Ala His Asn Leu Ser Val Gln Val Lys Lys Gly>

      700     710     720     730     740
      *      *      *      *      *
CCT TGG CAG ACT TTC TGT GCC TCT GAA TGG CCA ACA TTC GAT GTT GGA
Pro Trp Gln Thr Phe Cys Ala Ser Glu Trp Pro Thr Phe Asp Val Gly>

```

FIGURE 2

9/34

(SEQ ID NO: 2)
cont'd

```

      750      760      770      780      790
      *      *      *      *      *
TGG CCA TCA GAG GGG ACC TTT AAT TCT GAA ATT ATC CTG GCT GTT AAG
Trp Pro Ser Glu Gly Thr Phe Asn Ser Glu Ile Ile Leu Ala Val Lys>

      800      810      820      830      840
      *      *      *      *      *
GCA ATC ATT TTT CAG ACT GGA CCC GGC TCT CAT CCT GAT CAG GAG CCC
Ala Ile Ile Phe Gln Thr Gly Pro Gly Ser His Pro Asp Gln Glu Pro>

      850      860      870      880
      *      *      *      *
TAT ATC CTT ACG TGG CAA GAT TTG GCA GAA GAT CCT CCG CCA TGG GTT
Tyr Ile Leu Thr Trp Gln Asp Leu Ala Glu Asp Pro Pro Pro Trp Val>

      890      900      910      920      930
      *      *      *      *      *
AAA CCA TGG CTA AAT AAA CCA AGA AAG CCA GGT CCC CGA ATC CTG GCT
Lys Pro Trp Leu Asn Lys Pro Arg Lys Pro Gly Pro Arg Ile Leu Ala>

      940      950      960      970      980
      *      *      *      *      *
CTT GGA GAG AAA AAC AAA CAC TCG CCC GAA AAA GTC GAG CCC TCT CCT
Leu Gly Glu Lys Asn Lys His Ser Ala Glu Lys Val Glu Pro Ser Pro>

      990      1000      1010      1020      1030
      *      *      *      *      *
CGT ATC TAC CCC GAG ATC GAG GAG CCG CCG ACT TGG CCG GAA CCC CAA
Arg Ile Tyr Pro Glu Ile Glu Glu Pro Pro Thr Trp Pro Glu Pro Gln>

      1040      1050      1060      1070      1080
      *      *      *      *      *
CCT GTT CCC CCA CCC CCT TAT CCA GCA CAG GGT GCT GTG AGG GGA CCC
Pro Val Pro Pro Pro Pro Tyr Pro Ala Gln Gly Ala Val Arg Gly Pro>

      1090      1100      1110      1120
      *      *      *      *
TCT GCC CCT CCT GGA GCT CCG GTG GTG GAG GGA CCT OCT GCC GGG ACT
Ser Ala Pro Pro Gly Ala Pro Val Val Glu Gly Pro Ala Ala Gly Thr>

      1130      1140      1150      1160      1170
      *      *      *      *      *
CGG AGC CGG AGA GGC GCC ACC CCG GAG CCG ACA GAC GAG ATC GCG ATA
Arg Ser Arg Arg Gly Ala Thr Pro Glu Arg Thr Asp Glu Ile Ala Ile>

      1180      1190      1200      1210      1220
      *      *      *      *      *
TTA CCG CTG CCC ACC TAT GGC CCT CCC ATG CCA GGG GGC CAA TTG CAG
Leu Pro Leu Arg Thr Tyr Gly Pro Pro Met Pro Gly Gly Gln Leu Gln>

      1230      1240      1250      1260      1270
      *      *      *      *      *
CCC CTC CAG TAT TGG CCC TTT TCT TCT GCA GAT CTC TAT AAT TGG AAA
Pro Leu Gln Tyr Trp Pro Phe Ser Ser Ala Asp Leu Tyr Asn Trp Lys>

      1280      1290      1300      1310      1320
      *      *      *      *      *
ACT AAC CAT CCC CCT TTC TCG GAG GAT CCC CAA CCG CTC ACG GGG TTG
Thr Asn His Pro Pro Phe Ser Glu Asp Pro Gln Arg Leu Thr Gly Leu>

```

FIGURE 2, CONT.

10/34

(SEQ ID NO: 2)
cont'd

```

      1330      1340      1350      1360
      *      *      *      *
    GTG GAG TCC CTT ATG TTC TCT CAC CAG CCT ACT TGG GAT GAT TGT CAA
    Val Glu Ser Leu Met Phe Ser His Gln Pro Thr Trp Asp Asp Cys Gln>

1370      1380      1390      1400      1410
      *      *      *      *      *
    CAG CTG CTG CAG ACA CTC TTC ACA ACC GAG GAG CGA GAG AGA ATT CTG
    Gln Leu Leu Gln Thr Leu Phe Thr Thr Glu Glu Arg Glu Arg Ile Leu>

      1420      1430      1440      1450      1460
      *      *      *      *      *
    TTA GAG GCT AAA AAA AAT GTT CCT GGG GCC GAC GGG CGA CCC ACG CAG
    Leu Glu Ala Lys Lys Asn Val Pro Gly Ala Asp Gly Arg Pro Thr Gln>

      1470      1480      1490      1500      1510
      *      *      *      *      *
    TTG CAA AAT GAG ATT GAC ATG GGA TTT CCC TTG ACT CGC CCC GGT TGG
    Leu Cln Asn Glu Ile Asp Met Gly Phe Pro Leu Thr Arg Pro Gly Trp>

      1520      1530      1540      1550      1560
      *      *      *      *      *
    GAC TAC AAC ACG GCT GAA GGT ACG GAG AGC TTG AAA ATC TAT CCC CAG
    Asp Tyr Asn Thr Ala Glu Gly Arg Glu Ser Leu Lys Ile Tyr Arg Gln>

      1570      1580      1590      1600
      *      *      *      *
    GCT CTG GTG CGG GGT CTC CGG GGC GCC TCA AGA CGG CCC ACT AAT TTG
    Ala Leu Val Ala Gly Leu Arg Gly Ala Ser Arg Arg Pro Thr Asn Leu>

1610      1620      1630      1640      1650
      *      *      *      *      *
    GCT AAG GTA AGA GAG GTG ATG CAG GGA CCG AAC GAA CCT CCC TCG GTA
    Ala Lys Val Arg Glu Val Met Gln Gly Pro Asn Glu Pro Pro Ser Val>

      1660      1670      1680      1690      1700
      *      *      *      *      *
    TTT CTT GAG AGG CTC ATG GAA GCC TTC AGG CCG TTC ACC CCT TTT GAT
    Phe Leu Glu Arg Leu Met Glu Ala Phe Arg Arg Phe Thr Pro Phe Asp>

      1710      1720      1730      1740      1750
      *      *      *      *      *
    CCT ACC TCA GAG GCC CAG AAA GCC TCA GTG GCC CTG GCC TTC ATT GGG
    Pro Thr Ser Glu Ala Gln Lys Ala Ser Val Ala Leu Ala Phe Ile Gly>

      1760      1770      1780      1790      1800
      *      *      *      *      *
    CAG TCG GCT CTG GAT ATC AGG AAG AAA CTT CAG AGA CTG GAA GGG TTA
    Gln Ser Ala Leu Asp Ile Arg Lys Lys Leu Gln Arg Leu Glu Gly Leu>

      1810      1820      1830      1840
      *      *      *      *
    CAG GAG GCT GAG TTA CGT GAT CTA GTG AGA GAG GCA GAG AAG GTG TAT
    Gln Glu Ala Glu Leu Arg Asp Leu Val Arg Glu Ala Glu Lys Val Tyr>

1850      1860      1870      1880      1890
      *      *      *      *      *
    TAC AGA AGG GAG ACA GAA GAG GAG AAG GAA CAG AGA AAA GAA AAG GAG
    Tyr Arg Arg Glu Thr Glu Glu Glu Lys Glu Gln Arg Lys Glu Lys Glu>

```

FIGURE 2, CONT.

11/34

(SEQ ID NO: 2)
cont'd

```

1900      1910      1920      1930      1940
*         *         *         *         *
AGA GAA GAA ACG GAG GAA AGA CGT GAT AGA CCG CAA GAG AAG AAT TTG
Arg Glu Glu Arg Glu Glu Arg Arg Asp Arg Arg Gln Glu Lys Asn Leu>

1950      1960      1970      1980      1990
*         *         *         *         *
ACT AAG ATC TTG GCC GCA GTG GTT GAA GGG AAG AGC AGC AGG GAG AGA
Thr Lys Ile Leu Ala Ala Val Val Glu Gly Lys Ser Ser Arg Glu Arg>

2000      2010      2020      2030      2040
*         *         *         *         *
GAG AGA GAT TTT AGG AAA ATT AGG TCA GGC CCT AGA CAG TCA GGG AAC
Glu Arg Asp Phe Arg Lys Ile Arg Ser Gly Pro Arg Gln Ser Gly Asn>

2050      2060      2070      2080
*         *         *         *
CTG GGC AAT AGG ACC CCA CTC GAC AAG GAC CAG TGT GCG TAT TGT AAA
Leu Gly Asn Arg Thr Pro Leu Asp Lys Asp Gln Cys Ala Tyr Cys Lys>

2090      2100      2110      2120      2130
*         *         *         *         *
GAA AAA GGA CAC TGG GCA AGG AAC TGC CCC AAG AAG GGA AAC AAA GGA
Glu Lys Gly His Trp Ala Arg Asn Cys Pro Lys Lys Gly Asn Lys Gly>

2140      2150      2160      2170      2180
*         *         *         *         *
CCG AAG GTC CTA GCT CTA GAA GAA GAT AAA GAT T AGGGGAGACG
Pro Lys Val Leu Ala Leu Glu Glu Asp Lys Asp>

2190      2200      2210      2220      2230      2240
*         *         *         *         *         *
GGGTTCCGAC CCCCTCCCGC AGCCGAGGGT AACTTTGAAG GTGGAGGGGC AACCAGTTGA

2250      2260      2270      2280      2290      2300
*         *         *         *         *         *
GTTCTCTGGTT GATACCGGAG CGAGCATTC AGTGCTGCTA CAACCATTTAG GAAAACTAAA

2310      2320      2330      2340      2350
*         *         *         *         *
AGAAAAAATAA TCCTGGGTG ATG GGT GCC ACA GGG CAA CCG CAG TAT CCA TGG
Met Gly Ala Thr Gly Gln Arg Gln Tyr Pro Trp>

2360      2370      2380      2390      2400
*         *         *         *         *
ACT ACC CGA AGA ACC GTT GAC TTG CGA GTG GGA CCG GTA ACC CAC TCG
Thr Thr Arg Arg Thr Val Asp Leu Gly Val Gly Arg Val Thr His Ser>

2410      2420      2430      2440
*         *         *         *
TTT CTG GTC ATC CCT GAG TGC CCA GTA CCC CTT CTA GGT AGA GAC TTA
Phe Leu Val Ile Pro Glu Cys Pro Val Pro Leu Leu Gly Arg Asp Leu>

2450      2460      2470      2480      2490
*         *         *         *         *
CTG ACC AAG ATG GGA GCT CAA ATT TCT TTT GAA CAA CGA ACA CCA GAA
Leu Thr Lys Met Gly Ala Gln Ile Ser Phe Glu Gln Gly Arg Pro Glu>

```

FIGURE 2, CONT.

12/34

(SEQ ID NO: 2)
cont'd

```

2500      2510      2520      2530      2540
*         *         *         *         *
GTC TCT GTG AAT AAC AAA CCC ATC ACT GTG TTG ACC CTC CAA TTA GAT
Val Ser Val Asn Asn Lys Pro Ile Thr Val Leu Thr Leu Gln Leu Asp>

      2550      2560      2570      2580      2590
*         *         *         *         *
GAT GAA TAT CGA CTA TAT TCT CCC CAA GTA AAG CCT GAT CAA GAT ATA
Asp Glu Tyr Arg Leu Tyr Ser Pro Gln Val Lys Pro Asp Gln Asp Ile>

      2600      2610      2620      2630      2640
*         *         *         *         *
CAG TCC TCG TTG GAG CAG TTT CCC CAA GGC TGG GCA GAA ACC GCA GGG
Gln Ser Trp Leu Glu Gln Phe Pro Gln Ala Trp Ala Glu Thr Ala Gly>

      2650      2660      2670      2680
*         *         *         *
ATG GGT TTG CCA AAG CAA GTT CCC CCA CAG GTT ATT CAA CTG AAG GCC
Met Gly Leu Ala Lys Gln Val Pro Pro Gln Val Ile Gln Leu Lys Ala>

2690      2700      2710      2720      2730
*         *         *         *         *
AGT GCT ACA CCA GTA TCA GTC AGA CAG TAC CCC TTG AGT ACA GAG GCT
Ser Ala Thr Pro Val Ser Val Arg Gln Tyr Pro Leu Ser Arg Glu Ala>

      2740      2750      2760      2770      2780
*         *         *         *         *
CGA GAA GGA ATT TGG CCG CAT GTT CAA AGA TTA ATC CAA CAG GGC ATC
Arg Glu Gly Ile Trp Pro His Val Gln Arg Leu Ile Gln Gln Gly Ile>

      2790      2800      2810      2820      2830
*         *         *         *         *
CTA GTT CCT GTC CAA TCC CCT TGG AAT ACT CCC CTG CTA CCG GTT AGG
Leu Val Pro Val Gln Ser Pro Trp Asn Thr Pro Leu Leu Pro Val Arg>

      2840      2850      2860      2870      2880
*         *         *         *         *
AAG CCT GGG ACC AAT GAT TAT CGA CCA GTA CAG GAC TTG AGA GAG GTC
Lys Pro Gly Thr Asn Asp Tyr Arg Pro Val Gln Asp Leu Arg Glu Val>

      2890      2900      2910      2920
*         *         *         *
AAT AAA AGG GTG CAG GAC ATA CAC CCA ACG GTC CCG AAC CCT TAT AAC
Asn Lys Arg Val Gln Asp Ile His Pro Thr Val Pro Asn Pro Tyr Asn>

2930      2940      2950      2960      2970
*         *         *         *         *
CTC TTG AGC GCC CTC CCG CCT GAA CCG AAC TGG TAC ACA GTA TTG GAC
Leu Leu Ser Ala Leu Pro Pro Glu Arg Asn Trp Tyr Thr Val Leu Asp>

      2980      2990      3000      3010      3020
*         *         *         *         *
TTA AAA GAT GCC TTC TTC TGC CTG AGA TTA CAC CCC ACT AGC CAA CCA
Leu Lys Asp Ala Phe Phe Cys Leu Arg Leu His Pro Thr Ser Gln Pro>

      3030      3040      3050      3060      3070
*         *         *         *         *
CTT TTT ACC TTC GAA TGG AGA GAT CCA GGT ACG CGA AGA ACC GGG CAG
Leu Phe Thr Phe Glu Trp Arg Asp Pro Gly Thr Gly Arg Thr Gly Gln>

```

FIGURE 2. CONT.

13/34

(SEQ ID NO: 2)
cont'd

```

      3080      3090      3100      3110      3120
      *        *        *        *        *
CTC ACC TGG ACC CGA CTG CCC CAA GGG TTC AAG AAC TCC CCG ACC ATC
Leu Thr Trp Thr Arg Leu Pro Gln Gly Phe Lys Asn Ser Pro Thr Ile>

      3130      3140      3150      3160
      *        *        *        *
TTT GAC GAA GCC CTA CAC AGG GAC CTG GCC AAC TTC AGG ATC CAA CAC
Phe Asp Glu Ala Leu His Arg Asp Leu Ala Asn Phe Arg Ile Gln His>

3170      3180      3190      3200      3210
*        *        *        *        *
CCT CAG CTG ACC CTC CTC CAG TAC GTG GAT GAC CTG CTT CTG GCG CGA
Pro Gln Val Thr Leu Leu Gln Tyr Val Asp Asp Leu Leu Ala Gly>

      3220      3230      3240      3250      3260
      *        *        *        *        *
GCC ACC AAA CAG GAC TGC TTA GAA GGT ACG AAG GCA CTA CTG CTG GAA
Ala Thr Lys Gln Asp Cys Leu Glu Gly Thr Lys Ala Leu Leu Leu Glu>

      3270      3280      3290      3300      3310
      *        *        *        *        *
TTG TCT GAC CTA GGC TAC AGA GCC TCT GCT AAG AAG GCC CAG ATT TGC
Leu Ser Asp Leu Gly Tyr Arg Ala Ser Ala Lys Lys Ala Gln Ile Cys>

      3320      3330      3340      3350      3360
      *        *        *        *        *
AAG AGA GAG GTA ACA TAC TTG GGG TAC ACT TTG CCG GGC GGG CAG CGA
Arg Arg Glu Val Thr Tyr Leu Gly Tyr Ser Leu Arg Gly Gly Gln Arg>

      3370      3380      3390      3400
      *        *        *        *
TGG CTG ACG GAG GCA CGG AAG AAA ACT GTA CTC CAG ATA CCG GCC CCA
Trp Leu Thr Glu Ala Arg Lys Lys Thr Val Val Gln Ile Pro Ala Pro>

3410      3420      3430      3440      3450
*        *        *        *        *
ACC ACA GCC AAA CAA GTG AGA GAG TTT TTG GGG ACA GCT GGA TTT TGC
Thr Thr Ala Lys Gln Val Arg Glu Phe Leu Gly Thr Ala Gly Phe Cys>

      3460      3470      3480      3490      3500
      *        *        *        *        *
AGA CTG TGG ATC CCG GGG TTT CCG ACC TTA GCA GCC CCA CTC TAC CCG
Arg Leu Trp Ile Pro Gly Phe Ala Thr Leu Ala Ala Pro Leu Tyr Pro>

      3510      3520      3530      3540      3550
      *        *        *        *        *
CTA ACC AAA GAA AAA GGG GGT TGC TTA CCT CAG CAG GGA GGG AAA TA AAG
Leu Thr Lys Glu Lys Gly
      Lys Arg Gly Leu Leu Thr Ser Ala Gly Arg Glu Ile Lys>

      3560      3570      3580      3590      3600
      *        *        *        *        *
AAC AAA GAG GAA ATT CTA AGC CTA TTA GAA GCC TTA CAT TTG CCA AAA
Asn Lys Glu Glu Ile Leu Ser Leu Leu Glu Ala Leu His Leu Pro Lys>

      3610      3620      3630      3640      3650
      *        *        *        *        *
AGG CTA GCT ATT ATA CAC TGT CCT GGA CAT CAG AAA CCC AAA GAT CTC
Arg Leu Ala Ile Ile His Cys Pro Gly His Gln Lys Ala Lys Asp Leu>

```

FIGURE 2, CONT.

14/34

(SEQ ID NO: 2)
cont'd

```

      3660      3670      3680      3690
      *        *        *        *
ATA TCT AGA GGG AAC CAG ATG OCT GAC CGG GTT GCC AAG CAG GCA GCC
Ile Ser Arg Gly Asn Gln Met Ala Asp Arg Val Ala Lys Gln Ala Ala>

3700      3710      3720      3730      3740
      *        *        *        *        *
CAG GCT GTT AAC CTT CTG CCT ATA ATA GAA ACG CCC AAA GCC CCA GAA
Gln Ala Val Asn Leu Leu Pro Ile Ile Glu Thr Pro Lys Ala Pro Glu>

      3750      3760      3770      3780      3790
      *        *        *        *        *
CCC AGA CGA CAG TAC ACC CTA GAA GAC TGG CAA GAG ATA AAA AAG ATA
Pro Arg Arg Gln Tyr Thr Leu Glu Asp Trp Gln Glu Ile Lys Lys Ile>

      3800      3810      3820      3830      3840
      *        *        *        *        *
GAC CAG TTC TCT GAG ACT CCG GAG GGG ACC TGC TAT ACC TCA TAT GGG
Asp Gln Phe Ser Glu Thr Pro Glu Gly Thr Cys Tyr Thr Ser Tyr Gly>

      3850      3860      3870      3880      3890
      *        *        *        *        *
AAG GAA ATC CTG CCC CAC AAA GAA GGG TTA GAA TAT GTC CAA CAG ATA
Lys Glu Ile Leu Pro His Lys Glu Gly Leu Glu Tyr Val Gln Gln Ile>

      3900      3910      3920      3930
      *        *        *        *
CAT CGT CTA ACC CAC CTA GGA ACT AAA CAC CTG CAG CAG TTG GTC AGA
His Arg Leu Thr His Leu Gly Thr Lys His Leu Gln Gln Leu Val Arg>

3940      3950      3960      3970      3980
      *        *        *        *        *
ACA TCC CCT TAT CAT GTT CTG AGG CTA CCA GGA GTG CCT GAC TCG GTG
Thr Ser Pro Tyr His Val Leu Arg Leu Pro Gly Val Ala Asp Ser Val>

      3990      4000      4010      4020      4030
      *        *        *        *        *
GTC AAA CAT TGT GTG CCC TGC CAG CTG GTT AAT GAT AAT CCT TCC AGA
Val Lys His Cys Val Pro Cys Gln Leu Val Asn Ala Asn Pro Ser Arg>

      4040      4050      4060      4070      4080
      *        *        *        *        *
ATA CCT CCA GGA AAG AGA CTA AGG GGA AGC CAC CCA GGC GCT CAC TGG
Ile Pro Pro Gly Lys Arg Leu Arg Gly Ser His Pro Gly Ala His Trp>

      4090      4100      4110      4120      4130
      *        *        *        *        *
GAA GTG GAC TTC ACT GAG GTA AAG CCG GCT AAA TAC GGA AAC AAA TAT
Glu Val Asp Phe Thr Glu Val Lys Pro Ala Lys Tyr Gly Asn Lys Tyr>

      4140      4150      4160      4170
      *        *        *        *
CTA TTG GTT TTT GTA GAC ACC TTT TCA GGA TGG GTA GAG GCT TAT CCT
Leu Leu Val Phe Val Asp Thr Phe Ser Gly Trp Val Glu Ala Tyr Pro>

4180      4190      4200      4210      4220
      *        *        *        *        *
ACT AAA AAA GAG ACT TCA ACC GTG GTG CCT AAG AAA ATA CTG GAG GAA
Thr Lys Lys Glu Thr Ser Thr Val Val Ala Lys Lys Ile Leu Glu Glu>

```

FIGURE 2, CONT.

15/34

(SEQ ID NO: 2)
cont'd

```

4230      4240      4250      4260      4270
ATT TTT CCA AGA TTT OGA ATA CCT AAG GTA ATA GGG TCA GAC AAT GGT
Ile Phe Pro Arg Phe Gly Ile Pro Lys Val Ile Gly Ser Asp Asn Gly>

      4280      4290      4300      4310      4320
CCA GCT TTC GTT OCC CAG GTA ACT CAG GCA CTG GCC AAG ATA TTG GGG
Pro Ala Phe Val Ala Gln Val Ser Gln Gly Leu Ala Lys Ile Leu Gly>

      4330      4340      4350      4360      4370      4380
ATT GAT TG A AAA CTG CAT TGT GCA TAC AGA CCC CAA AGC TCA OGA CAG
Ile Asp      Lys Leu His Cys Ala Tyr Arg Pro Gln Ser Ser Gly Gln>

      4380      4390      4400      4410
GTA GAG AGG ATG AAT AGA ACC ATT AAA GAG ACC CTT ACC AAA TTG ACC
Val Glu Arg Met Asn Arg Thr Ile Lys Glu Thr Leu Thr Lys Leu Thr>

4420      4430      4440      4450      4460
ACA GAG ACT GGC ATT AAT GAT TGG ATG GCT CTC CTG CCC TTT GTG CTT
Thr Glu Thr Gly Ile Asn Asp Trp Met Ala Leu Leu Pro Phe Val Leu>

      4470      4480      4490      4500      4510
TTT AGG GTG AGG AAC ACC CCT GGA CAG TTT GGG CTG ACC CCC TAT AAA
Phe Arg Val Arg Asn Thr Pro Gly Gln Phe Gly Leu Thr Pro Tyr Lys>

      4520      4530      4540      4550      4560
TTG CTC TAC GGG GGA CCC CCC CCG TTG GCA GAA ATT GCC TTT GCA CAT
Leu Leu Tyr Gly Gly Pro Pro Pro Leu Ala Glu Ile Ala Phe Ala His>

      4570      4580      4590      4600      4610
AGT GCT GAT GTG CTG CTT TCC CAG CCT TTG TTC TCT ACG CTC AAG GCG
Ser Ala Asp Val Leu Leu Ser Gln Pro Leu Phe Ser Arg Leu Lys Ala>

      4620      4630      4640      4650
CTC GAG TGG GTG ACG CAG CGA GCG TGG AAG CAG CTC CCG GAG GCC TAC
Leu Glu Trp Val Arg Gln Arg Ala Trp Lys Gln Leu Arg Glu Ala Tyr>

4660      4670      4680      4690      4700
TCA OGA OGA CAC TTG CAA GTT CCA CAT CGC TTC CAA GTT GGA GAT TCA
Ser Gly Gly Asp Leu Gln Val Pro His Arg Phe Gln Val Gly Asp Ser>

      4710      4720      4730      4740      4750
GTC TAT GTT AGA CGC CAC CGT GCA GGA AAC CTC CAG ACT CCG TAG AAG
Val Tyr Val Arg Arg His Arg Ala Gly Asn Leu Glu Thr Arg *** Lys>

      4760      4770      4780      4790      4800
OGA CCT TAT CTC GTA CTT TTG ACC ACA CCA ACG GCT GTG AAA GTC GAA
Gly Pro Tyr Leu Val Leu Leu Thr Thr Pro Thr Ala Val Lys Val Glu>

```

FIGURE 2, CONT.

16/34

(SEQ ID NO: 2)
cont'd

```

      4810      4820      4830      4840      4850
      *         *         *         *         *
GGA ATC CCC TTA AGC TTC GCC TCC ATC GCG TGG TTC CTT ACT CTG TCA
Gly Ile Pro Leu Ser Phe Ala Ser Ile Ala Trp Phe Leu Thr Leu Ser>

      4860      4870      4880      4890
      *         *         *         *
ATA ACT CCT CAA GTT AAT GGT AAA CCG CTT GTG GAC AGC CCG AAC TCC
Ile Thr Pro Gln Val Asn Gly Lys Arg Leu Val Asp Ser Pro Asn Ser>

4900      4910      4920      4930      4940
      *         *         *         *         *
CAT AAA CCC TTA TCT CTC ACC TGG TTA CTT ACT GAC TCC GGT ACA GGT
His Lys Pro Leu Ser Leu Thr Trp Leu Leu Thr Asp Ser Gly Thr Gly>

      4950      4960      4970      4980      4990
      *         *         *         *         *
ATT AAT ATT AAC AGC ACT CAA CCG GAG GCT CCC TTG GGG ACC TGG TGG
Ile Asn Ile Asn Ser Thr Gln Gly Glu Ala Pro Leu Gly Thr Trp Trp>

      5000      5010      5020      5030      5040
      *         *         *         *         *
CCT GAA TTA TAT GTC TGC CTT CGA TCA GTA ATC CCT GGT CTC AAT GAC
Pro Glu Leu Tyr Val Cys Leu Arg Ser Val Ile Pro Gly Leu Asn Asp>

      5050      5060      5070      5080      5090
      *         *         *         *         *
CAG GCC ACA CCC CCC GAT GTA CTC GGT GCT TAC GGG TTT TAC GTT TGC
Gln Ala Thr Pro Pro Asp Val Leu Arg Ala Tyr Gly Phe Tyr Val Cys>

      5100      5110      5120      5130
      *         *         *         *
CCA GGA CCC CCA AAT AAT GAA GAA TAT TGT GGA AAT CCT CAG GAT TTC
Pro Gly Pro Pro Asn Asn Glu Glu Tyr Cys Gly Asn Pro Gln Asp Phe>

5140      5150      5160      5170      5180
      *         *         *         *         *
TTT TGC AAG CAA TGG AGC TGC ATA ACT TCT AAT GAT GGG AAT TGG AAA
Phe Cys Lys Gln Trp Ser Cys Ile Thr Ser Asn Asp Gly Asn Trp Lys>

      5190      5200      5210      5220      5230
      *         *         *         *         *
TOG CCA GTC TCT CAG CAA GAC AGA GTA AGT TAC TCT TTT GTT AAC AAT
Trp Pro Val Ser Gln Gln Asp Arg Val Ser Tyr Ser Phe Val Asn Asn>

      5240      5250      5260      5270      5280
      *         *         *         *         *
CCT ACC AGT TAT AAT CAA TTT AAT TAT GCG CAT GCG AGA TGG AAA GAT
Pro Thr Ser Tyr Asn Gln Phe Asn Tyr Gly His Gly Arg Trp Lys Asp>

      5290      5300      5310      5320      5330
      *         *         *         *         *
TOG CAA CAG CCG GTA CAA AAA GAT GTA CGA AAT AAG CAA ATA AGC TGT
Trp Gln Gln Arg Val Gln Lys Asp Val Arg Asn Lys Gln Ile Ser Cys>

      5340      5350      5360      5370
      *         *         *         *
CAT TCG TTA GAC CTA GAT TAC TTA AAA ATA AGT TTC ACT GAA AAA GGA
His Ser Leu Asp Leu Asp Tyr Leu Lys Ile Ser Phe Thr Glu Lys Gly>

```

FIGURE 2, CONT.

17/34

(SEQ ID NO: 2)
cont'd

5380 5390 5400 5410 5420
* * * * *
AAA CAA GAA AAT ATT CAA AAG TGG GTA AAT GGT ATA TCT TGG GGA ATA
Lys Gln Glu Asn Ile Gln Lys Trp Val Asn Gly Ile Ser Trp Gly Ile>

5430 5440 5450 5460 5470
* * * * *
GTG TAC TAT GGA OCC TCT GGG AGA AAG AAA GGA TCT GTT CTG ACT ATT
Val Tyr Tyr Gly Gly Ser Gly Arg Lys Lys Gly Ser Val Leu Thr Ile>

5480 5490 5500 5510 5520
* * * * *
CCG CTC AGA ATA GAA ACT CAG ATG GAA CCT CCG GTT GCT ATA GGA CCA
Arg Leu Arg Ile Glu Thr Gln Met Glu Pro Pro Val Ala Ile Gly Pro>

5530 5540 5550 5560
* * * *
AAT AAG GGT TTG GCC GAA CAA GGA CCT CCA ATC CAA GAA CAG
Asn Lys Gly Leu Ala Glu Gln Gly Pro Pro Ile Gln Glu Gln>

5570 5580 5590 5600 5610
* * * * *
AGG CCA TCT CCT AAC CCC TCT GAT TAC AAT ACA ACC TCT GGA TCA GTC
Arg Pro Ser Pro Asn Pro Ser Asp Tyr Asn Thr Thr Ser Gly Ser Val>

5620 5630 5640 5650 5660
* * * * *
CCC ACT GAG CCT AAC ATC ACT ATT AAA ACA GGG GCG AAA CTT TTT AGC
Pro Thr Glu Pro Asn Ile Thr Ile Lys Thr Gly Ala Lys Leu Phe Ser>

5670 5680 5690 5700
* * * *
CTC ATC CAG GGA GCT TTT CAA GCT CTT AAC TCC ACG ACT CCA GAG GCT
Leu Ile Gln Gly Ala Phe Gln Ala Leu Asn Ser Thr Thr Pro Glu Ala>

5710 5720 5730 5740 5750
* * * * *
ACC TCT TCT TGT TGG CTT TGC TTA GCT TCG GGC CCA CCT TAC TAT GAG
Thr Ser Ser Cys Trp Leu Cys Leu Ala Ser Gly Pro Pro Tyr Tyr Glu>

5760 5770 5780 5790 5800
* * * * *
GGA ATG OCT AGA GGA GGG AAA TTC AAT GTG ACA AAG GAA CAT AGA GAC
Gly Met Ala Arg Gly Gly Lys Phe Asn Val Thr Lys Glu His Arg Asp>

5810 5820 5830 5840 5850
* * * * *
CAA TGT ACA TGG GGA TCC CAA AAT AAG CTT ACC CTT ACT GAG GTT TCT
Gln Cys Thr Trp Gly Ser Gln Asn Lys Leu Thr Leu Thr Glu Val Ser>

5860 5870 5880 5890 5900
* * * * *
GGA AAA GGC ACC TGC ATA GGG ATG GTT CCC CCA TCC CAC CAA CAC CTT
Gly Lys Gly Thr Cys Ile Gly Met Val Pro Pro Ser His Gln His Leu>

5910 5920 5930 5940
* * * *
TGT AAC CAC ACT GAA GCC TTT AAT CGA ACC TCT GAG AGT CAA TAT CTG
Cys Asn His Thr Glu Ala Phe Asn Arg Thr Ser Glu Ser Gln Tyr Leu>

FIGURE 2. CONT.

18/34

(SEQ ID NO: 2)
cont'd

```

5950      5960      5970      5980      5990
*         *         *         *         *
GTA CCT GGT TAT GAC AGG TGG TGG GCA TGT AAT ACT GGA TTA ACC CCT
Val Pro Gly Tyr Asp Arg Trp Trp Ala Cys Asn Thr Gly Leu Thr Pro>

6000      6010      6020      6030      6040
*         *         *         *         *
TGT GTT TCC ACC TTG GTT TTC AAC CAA ACT AAA GAC TTT TGC GTT ATG
Cys Val Ser Thr Leu Val Phe Asn Gln Thr Lys Asp Phe Cys Val Met>

6050      6060      6070      6080      6090
*         *         *         *         *
GTC CAA ATT GTC CCC CGG GTG TAC TAC TAT CCC GAA AAA GCA GTC CTT
Val Gln Ile Val Pro Arg Val Tyr Tyr Tyr Pro Glu Lys Ala Val Leu>

6100      6110      6120      6130      6140
*         *         *         *         *
GAT GAA TAT GAC TAT AGA TAT AAT CGG CCA AAA AGA GAG CCC ATA TCC
Asp Glu Tyr Asp Tyr Arg Tyr Asn Arg Pro Lys Arg Glu Pro Ile Ser>

6150      6160      6170      6180
*         *         *         *
CTG ACA CTA GCT GTA ATG CTC GGA TTG GGA GTG GCT GCA GCC GTG GGA
Leu Thr Leu Ala Val Met Leu Gly Leu Gly Val Ala Ala Gly Val Gly>

6190      6200      6210      6220      6230
*         *         *         *         *
ACA GGA ACG GCT GCC CTA ATC ACA GGA CCG CAA CAG CTG GAG AAA GGA
Thr Gly Thr Ala Ala Leu Ile Thr Gly Pro Gln Gln Leu Glu Lys Gly>

6240      6250      6260      6270      6280
*         *         *         *         *
CTT AGT AAC CTA CAT CGA ATT GTA ACG GAA GAT CTC CAA GCC CTA GAA
Leu Ser Asn Leu His Arg Ile Val Thr Glu Asp Leu Gln Ala Leu Glu>

6290      6300      6310      6320      6330
*         *         *         *         *
AAA TCT GTC AGT AAC CTG GAG GAA TCC CTA ACC TCC TTA TCT GAA GTG
Lys Ser Val Ser Asn Leu Glu Glu Ser Leu Thr Ser Leu Ser Glu Val>

6340      6350      6360      6370      6380
*         *         *         *         *
GTT CTA CAG AAC AGA AGG GGG TTA GAT CTG TTA TTT CTA AAA GAA GGA
Val Leu Gln Asn Arg Arg Gly Leu Asp Leu Leu Phe Leu Lys Glu Gly>

6390      6400      6410      6420
*         *         *         *
GGG TTA TGT GTA GCC TTA AAA GAG GAA TGC TGC TTC TAT GTA GAT CAC
Gly Leu Cys Val Ala Leu Lys Glu Glu Cys Cys Phe Tyr Val Asp His>

6430      6440      6450      6460      6470
*         *         *         *         *
TCA GGA GCC ATC AGA GAC TCC ATG AGC AAG CTT AGA GAA AGG TTA GAG
Ser Gly Ala Ile Arg Asp Ser Met Ser Lys Leu Arg Glu Arg Leu Glu>

6480      6490      6500      6510      6520
*         *         *         *         *
AGG CGT CGA AGG GAA AGA GAG GCT GAC CAG GGG TGG TTT GAA GGA TGG
Arg Arg Arg Arg Glu Arg Glu Ala Asp Gln Gly Trp Phe Glu Gly Trp>

```

FIGURE 2, CONT.

19/34

(SEQ ID NO: 2)
cont'd

6530 6540 6550 6560 6570
* * * * *
TTC AAC AGG TCT CCT TGG ATG ACC ACC CTG CTT TCT GCT CTG ACC GGG
Phe Asn Arg Ser Pro Trp Met Thr Thr Leu Leu Ser Ala Leu Thr Gly>

6580 6590 6600 6610 6620
* * * * *
CCC CTA GTA GTC CTG CTC CTG TTA CTT ACA GTT GCG CCT TGC TTA ATT
Pro Leu Val Val Leu Leu Leu Leu Leu Thr Val Gly Pro Cys Leu Ile>

6630 6640 6650 6660
* * * * *
AAT AGG TTT GTT GGC TTT GTT AGA GAA CGA GTG AGT GCA GTC CAG ATC
Asn Arg Phe Val Ala Phe Val Arg Glu Arg Val Ser Ala Val Gln Ile>

6670 6680 6690 6700 6710
* * * * *
ATG GTA CTT AGG CAA CAG TAC CAA GGC CTT CTG AGC CAA GGA GAA ACT
Met Val Leu Arg Gln Gln Tyr Gln Gly Leu Leu Ser Gln Gly Glu Thr>

6720 6730 6740 6750 6760 6770
* * * * *
GAC CTC TAGCCTTC CCACTTCTAA GATTAGAACT ATTAACAAGA CAAGAAGTGG
Asp Leu>

6780 6790 6800 6810 6820 6830
* * * * *
OGAATGAAAG GATGAAATG CAACCTAACC CTCCAGAAC CCAGGAAGTT AATAAAAAGC

6840 6850 6860 6870 6880 6890
* * * * *
TCTAAATGCC CCGAATTC AGACCTGCT GCGTCCAGT AAATAGGTAG AAGGTACAC

6900 6910 6920 6930 6940 6950
* * * * *
TTCTATTGT TCCAGGGCCT GCTATCCTGG CCTAAGTAAG ATAACAGGAA ATGAGTTGAC

6960 6970 6980 6990 7000 7010
* * * * *
TAATCCCTTA TCTGGATTCT GTAAACTGA CTGGCACCAT AGAAGAATTG ATTACACATT

7020 7030 7040 7050 7060 7070
* * * * *
GACAGCCCTA GTGACCTATC TCAACTGCAA TCTGTCACTC TGCCAGGAG CCCACOCAGA

7080 7090 7100 7110 7120 7130
* * * * *
TGCGGACCTC CGGAGCTATT TTAAATGAT TGGTCCACGG AGCGGGGCT CTCGATATTT

7140 7150 7160 7170 7180 7190
* * * * *
TAAAAAGATT GGTCCATGGA GCGCGGGCTC TCGATATTTT AAAATGATTG GPTGTGACG

7200 7210 7220 7230 7240 7250
* * * * *
CACAGGCTTT GTGTGAACC CCATAAAAGC TGTCCGATT CCGCACTGG GCGCGCACTC

FIGURE 2, CONT.

20/34

7260 7270 7280 7290 7300 7310 (SEQ ID NO: 2)
* * * * * cont'd
CTCTACCCCT GUGTGGTGTA CGACTGTGCG CCCACGCGG CTTGGAATAA AAATCCTCTT
7320 7330
* * * * *
GCTGTTTGCA TCAAAAAAAAA AAA

FIGURE 2, CONT.

SUBSTITUTE SHEET (RULE 26)

21/34

(SEQ ID NO: 3)

```

      10      20      30      40      50      60
      *      *      *      *      *      *
CCGTGCTGTA CGACTGTGGG CCCAGCGCG CTGGGAATAA AAATCCTCTT GCTGTTTGCA

      70      80      90     100     110     120
      *      *      *      *      *      *
TCAAGACCGC TTCTCGTGAG TGATTAAGGG GAGTCGCCCT TCCGAGCCT GGAGGTTCTT

     130     140     150     160     170     180
      *      *      *      *      *      *
TTTGCTGGTC TTACATTGGG GGGCTGTGCC GGGATCTGTC GCGGCCACCC CTAACACCCG

     190     200     210     220     230     240
      *      *      *      *      *      *
AGAAACGACT TGGAGGTAAA AAGGATCCTC TTTTAAAGT GTATGCATGT ACCGCGCGGC

     250     260     270     280     290     300
      *      *      *      *      *      *
GTCTCTGTTT TGAGTGTCTG TTTTCAGTGG TCGCGCCTT CGTTTTCAG CTGTCTCTC

     310     320     330     340     350     360
      *      *      *      *      *      *
AGGCCGTAAG GGCTGGGGGA CTGTGATCAG CAGAGCTGCT AGGAAGATCA CAGGCTGCTG

     370     380     390     400     410     420
      *      *      *      *      *      *
CCCTGGGGGA CCGCCCGGGA GGTGAAGGGA OCCAGGACG CCTGGTGGTC TCCTACTGTC

     430     440     450     460     470     480
      *      *      *      *      *      *
GGTCAGAGGA CCGAATTCTG TTGCTGAAGC GAAAGCTTCC CCCTCCGGGA CCGTCCGACT

     490     500     510     520     530     540
      *      *      *      *      *      *
CTTTGCGCTG CTGTGGAAG ACGTGACCG GTACCGTGTG TCTGGATCTG TTGGTTTCTG

     550     560     570     580     590
      *      *      *      *      *
TTTGTGTGT CTTTGTCTTG TGTCGCTTG TCTACAGTTT TAAT ATG GGA CAG ACG
                                   Met Gly Gln Thr>

     600     610     620     630     640
      *      *      *      *      *
GTG ACG ACC CCT CTT AGT TTG ACT CTC GAC CAT TGG ACT GAA GTT AAA
Val Thr Thr Pro Leu Ser Leu Thr Leu Asp His Trp Thr Glu Val Lys>

     650     660     670     680     690
      *      *      *      *      *
TCC AGG CCT CAT AAT TTG TCA GTT CAG GTT AAG AAG GGA CCT TGG CAG
Ser Arg Ala His Asn Leu Ser Val Gln Val Lys Lys Gly Pro Trp Gln>

     700     710     720     730     740
      *      *      *      *      *
ACT TTC TGT GTC TCT GAA TGG CCG ACA TTC GAT GTT GGA TGG CCA TCA
Thr Phe Cys Val Ser Glu Trp Pro Thr Phe Asp Val Gly Trp Pro Ser>

```

FIGURE 3

22/34

(SEQ ID NO: 3)
cont'd

```

      750      760      770      780
      *      *      *      *
GAG GGG ACC TTT AAT TCT GAG ATT ATC CTG GCT GTT AAA GCA GTT ATT
Glu Gly Thr Phe Asn Ser Glu Ile Ile Leu Ala Val Lys Ala Val Ile>

790      800      810      820      830
      *      *      *      *      *
TTT CAG ACT GGA CCC GGC TCT CAT CCC GAT CAG GAG CCC TAT ATC CTT
Phe Gln Thr Gly Pro Gly Ser His Pro Asp Gln Glu Pro Tyr Ile Leu>

      840      850      860      870      880
      *      *      *      *      *
ACG TGG CAA GAT TTG GCA GAG GAT CCT CCG CCA TGG GTT AAA CCA TGG
Thr Trp Gln Asp Leu Ala Glu Asp Pro Pro Pro Trp Val Lys Pro Trp>

      890      900      910      920      930
      *      *      *      *      *
CTG AAT AAG CCA AGA AAG CCA GGT CCC CGA ATT CTG GCT CTT GGA GAG
Leu Asn Lys Pro Arg Lys Pro Gly Pro Arg Ile Leu Ala Leu Gly Glu>

      940      950      960      970      980
      *      *      *      *      *
AAA AAC AAA CAC TCG CCT GAA AAA GTC AAG CCC TCT CCT CAT ATC TAC
Lys Asn Lys His Ser Ala Glu Lys Val Lys Pro Ser Pro His Ile Tyr>

      990      1000      1010      1020
      *      *      *      *
CCC GAG ATT GAG GAG CCA CCG GCT TGG CCG GAA CCC CAA TCT GTT CCC
Pro Glu Ile Glu Glu Pro Pro Ala Trp Pro Glu Pro Gln Ser Val Pro>

1030      1040      1050      1060      1070
      *      *      *      *      *
CCA CCC CCT TAT CTG GCA CAG GGT GCC GCG AGG CGA CCC TTT GCC CCT
Pro Pro Pro Tyr Leu Ala Gln Gly Ala Ala Arg Gly Pro Phe Ala Pro>

      1080      1090      1100      1110      1120
      *      *      *      *      *
CCT GGA GCT CCG GCG GTG GAG CGA CCT GCT GCA GCG ACT CCG AGC CCG
Pro Gly Ala Pro Ala Val Glu Gly Pro Ala Ala Gly Thr Arg Ser Arg>

      1130      1140      1150      1160      1170
      *      *      *      *      *
AGG GGC GCC ACC CCG GAG CCG ACA GAC GAG ATC GCG ACA TTA CCG CTG
Arg Gly Ala Thr Pro Glu Arg Thr Asp Glu Ile Ala Thr Leu Pro Leu>

      1180      1190      1200      1210      1220
      *      *      *      *      *
CGC ACG TAC GGC CCT CCC ACA CCG GCG GGC CAA TTG CAG CCC CTC CAG
Arg Thr Tyr Gly Pro Pro Thr Pro Gly Gly Gln Leu Gln Pro Leu Gln>

      1230      1240      1250      1260
      *      *      *      *
TAT TGG CCC TTT TCT TCT GCA GAT CTC TAT AAT TGG AAA ACT AAC CAT
Tyr Trp Pro Phe Ser Ser Ala Asp Leu Tyr Asn Trp Lys Thr Asn His>

1270      1280      1290      1300      1310
      *      *      *      *      *
CCC CCT TTC TCG GAG GAT CCC CAA CCG CTC ACG GCG TTG GTG GAG TCC
Pro Pro Phe Ser Glu Asp Pro Gln Arg Leu Thr Gly Leu Val Glu Ser>

```

FIGURE 3,CONT.

23/34

(SEQ ID NO: 3)
cont'd

```

1320      1330      1340      1350      1360
*         *         *         *         *
CTT ATG TTC TCT CAC CAG CCT ACT TGG GAT GAT TGT CAA CAG CTG CTG
Leu Met Phe Ser His Gln Pro Thr Trp Asp Asp Cys Gln Gln Leu Leu>

1370      1380      1390      1400      1410
*         *         *         *         *
CAG ACA CTC TTC ACA ACC GAG GAG CGA GAG AGA ATT CTA TTA GAG GCT
Gln Thr Leu Phe Thr Thr Glu Glu Arg Glu Arg Ile Leu Leu Glu Ala>

1420      1430      1440      1450      1460
*         *         *         *         *
AGA AAA AAT GTT CCT GGG GCC GAC GGG CGA CCC ACG CGG TTG CAA AAT
Arg Lys Asn Val Pro Gly Ala Asp Gly Arg Pro Thr Arg Leu Gln Asn>

1470      1480      1490      1500
*         *         *         *
GAG ATT GAC ATG GGA TTT CCC TTA ACT CGC CCC GGT TGG GAC TAC AAC
Glu Ile Asp Met Gly Phe Pro Leu Thr Arg Pro Gly Trp Asp Tyr Asn>

1510      1520      1530      1540      1550
*         *         *         *         *
ACG GCT GAA GGT AGG GAG AGC TTG AAA ATC TAT CGC CAG GCT CTG GTG
Thr Ala Glu Gly Arg Glu Ser Leu Lys Ile Tyr Arg Gln Ala Leu Val>

1560      1570      1580      1590      1600
*         *         *         *         *
GGG GGT CTC CGG GGC GCC TCA AGA CGG CCC ACT AAT TTG OCT AAG GTA
Ala Gly Leu Arg Gly Ala Ser Arg Arg Pro Thr Asn Leu Ala Lys Val>

1610      1620      1630      1640      1650
*         *         *         *         *
AGA GAA GTG ATG CAG GGA CGG AAT GAA CCC CCC TCT GTT TTT CTT GAG
Arg Glu Val Met Gln Gly Pro Asn Glu Pro Pro Ser Val Phe Leu Glu>

1660      1670      1680      1690      1700
*         *         *         *         *
AGG CTC TTG GAA GCC TTC AGG CGG TAC ACC CCT TTT GAT CCC ACC TCA
Arg Leu Leu Glu Ala Phe Arg Arg Tyr Thr Pro Phe Asp Pro Thr Ser>

1710      1720      1730      1740
*         *         *         *
GAG GCC CAA AAA GCC TCA GTG GCT TTG GCC TTT ATA GGA CAG TCA GCC
Glu Ala Gln Lys Ala Ser Val Ala Leu Ala Phe Ile Gly Gln Ser Ala>

1750      1760      1770      1780      1790
*         *         *         *         *
TTG GAT ATT AGA AAG AAG CTT CAG AGA CTG GAA GGG TTA CAG GAG GCT
Leu Asp Ile Arg Lys Lys Leu Gln Arg Leu Glu Gly Leu Gln Glu Ala>

1800      1810      1820      1830      1840
*         *         *         *         *
GAG TTA CGT GAT CTA GTG AAG GAG GCA GAG AAA GTA TAT TAC AAA AGG
Glu Leu Arg Asp Leu Val Lys Glu Ala Glu Lys Val Tyr Tyr Lys Arg>

1850      1860      1870      1880      1890
*         *         *         *         *
GAG ACA GAA GAA GAA AGG GAA CAA AGA AAA GAG AGA GAA AGA GAG GAA
Glu Thr Glu Glu Glu Arg Glu Gln Arg Lys Glu Arg Glu Arg Glu Glu>

```

FIGURE 3,CONT.

24/34

(SEQ ID NO: 3)
cont'd

```

      1900      1910      1920      1930      1940
      *         *         *         *         *
AGG GAG GAA AGA CGT AAT AAA CCG CAA GAG AAG AAT TTG ACT AAG ATC
Arg Glu Glu Arg Arg Asn Lys Arg Gln Glu Lys Asn Leu Thr Lys Ile>

      1950      1960      1970      1980
      *         *         *         *         *
TTG GCT GCA GTG GTT GAA GGG AAA AGC AAT ACG GAA AGA GAG AGA GAT
Leu Ala Ala Val Val Glu Gly Lys Ser Asn Thr Glu Arg Glu Arg Asp>

1990      2000      2010      2020      2030
      *         *         *         *         *
TTT AGG AAA ATT AGG TCA GGC CCT AGA CAG TCA GGG AAC CTG GGC AAT
Phe Arg Lys Ile Arg Ser Gly Pro Arg Gln Ser Gly Asn Leu Gly Asn>

      2040      2050      2060      2070      2080
      *         *         *         *         *
AGG ACC CCA CTC GAC AAG GAC CAA TGT GCA TAT TGT AAA GAA AGA GGA
Arg Thr Pro Leu Asp Lys Asp Gln Cys Ala Tyr Cys Lys Glu Arg Gly>

      2090      2100      2110      2120      2130
      *         *         *         *         *
CAC TGG GCA AGG AAC TGC CCC AAG AAG GGA AAC AAA GGA CCA AGG ATC
His Trp Ala Arg Asn Cys Pro Lys Lys Gly Asn Lys Gly Pro Arg Ile>

      2140      2150      2160      2170      2180
      *         *         *         *         *
CTA GCT CTA GAA GAA GAT AAA GAT TACG GGAGACGGGG TTCGACCCCC
Leu Ala Leu Glu Glu Asp Lys Asp>

      2190      2200      2210      2220      2230      2240
      *         *         *         *         *         *
CTCCCGAGC CCAGGTAAC TTTGAAGGTG GAGGGCAAC CACTTGAGTT CCTGGTTGAT

      2250      2260      2270      2280      2290      2300
      *         *         *         *         *         *
ACCGAGCGA AACATTCACT GCTACTACAG CCATTAGGAA AACTAAAAGA TAAAAAATCC

      2310      2320      2330      2340      2350
      *         *         *         *         *
TGGGTG ATG GGT GCC ACA GGG CAA CAA CAG TAT CCA TGG ACT ACC CGA AGA
Met Gly Ala Thr Gly Gln Gln Gln Tyr Pro Trp Thr Thr Arg Arg>

      2360      2370      2380      2390
      *         *         *         *         *
ACA GTT GAC TTG GGA GTG GGA CGG GTA ACC CAC TCG TTT CTG GTC ATA
Thr Val Asp Leu Gly Val Gly Arg Val Thr His Ser Phe Leu Val Ile>

2400      2410      2420      2430      2440
      *         *         *         *         *
CCT GAG TGC CCA GCA CCC CTC TTA GGT AGA GAC TTA TTG ACC AAG ATG
Pro Glu Cys Pro Ala Pro Leu Leu Gly Arg Asp Leu Leu Thr Lys Met>

2450      2460      2470      2480      2490
      *         *         *         *         *
GGA GCA CAA ATT TCT TTT GAA CAA GGG AAA CCA GAA GTG TCT GCA AAT
Gly Ala Gln Ile Ser Phe Glu Gln Gly Lys Pro Glu Val Ser Ala Asn>

```

FIGURE 3,CONT.

25/34

(SEQ ID NO: 3)
cont'd

```

      2500      2510      2520      2530      2540
      *         *         *         *         *
AAC AAA OCT ATC ACT GTG TTG ACC CTC CAA TTA GAT GAC GAA TAT CGA
Asn Lys Pro Ile Thr Val Leu Thr Leu Gln Leu Asp Asp Glu Tyr Arg>

      2550      2560      2570      2580      2590
      *         *         *         *         *
CTA TAC TCT CCC CTA GTA AAG CCT GAT CAA AAT ATA CAA TTC TGG TTG
Leu Tyr Ser Pro Leu Val Lys Pro Asp Gln Asn Ile Gln Phe Trp Leu>

      2600      2610      2620      2630
      *         *         *         *
GAA CAG TTT CCC CAA GCC TGG GCA GAA ACC GCA GGG ATG GGT TTG GCA
Glu Gln Phe Pro Gln Ala Trp Ala Glu Thr Ala Gly Met Gly Leu Ala>

2640      2650      2660      2670      2680
      *         *         *         *         *
AAG CAA GTT CCC CCA CAA GTT ATT CAA CTG AAG GCC AGT GCC ACA CCA
Lys Gln Val Pro Pro Gln Val Ile Gln Leu Lys Ala Ser Ala Thr Pro>

2690      2700      2710      2720      2730
      *         *         *         *         *
GTG TCA CTC AGA CAG TAC CCC TTG AGT AAA GAA GCT CAA GAA GGA ATT
Val Ser Val Arg Gln Tyr Pro Leu Ser Lys Glu Ala Gln Glu Gly Ile>

      2740      2750      2760      2770      2780
      *         *         *         *         *
CGG CCG CAT GTC CAA CCA TTA ATC CAA CAG GCC ATC CTA GTT CCT GTC
Arg Pro His Val Gln Arg Leu Ile Gln Gln Gly Ile Leu Val Pro Val>

      2790      2800      2810      2820      2830
      *         *         *         *         *
CAA TCT CCC TGG AAT ACT CCC CTG CTA CCG GTT AGA AAG CCT GGG ACT
Gln Ser Pro Trp Asn Thr Pro Leu Leu Pro Val Arg Lys Pro Gly Thr>

      2840      2850      2860      2870
      *         *         *         *
AAT GAC TAT CGA CCA GTA CAG GAC TTG AGA GAG GTC AAT AAA CCG GIG
Asn Asp Tyr Arg Pro Val Gln Asp Leu Arg Glu Val Asn Lys Arg Val>

2880      2890      2900      2910      2920
      *         *         *         *         *
CAG GAT ATA CAC CCA ACA GTC CCG AAC CCT TAT AAC CTC TTG TGT GCT
Gln Asp Ile His Pro Thr Val Pro Asn Pro Tyr Asn Leu Leu Cys Ala>

2930      2940      2950      2960      2970
      *         *         *         *         *
CTC CCA CCC CAA CCG ACC TGG TAT ACA GTA TTG GAC TTA AAG GAT GCC
Leu Pro Pro Gln Arg Ser Trp Tyr Thr Val Leu Asp Leu Lys Asp Ala>

      2980      2990      3000      3010      3020
      *         *         *         *         *
TTC TTC TGC CTG AGA TTA CAC CCC ACT AGC CAA CCA CTT TTT GCC TTC
Phe Phe Cys Leu Arg Leu His Pro Thr Ser Gln Pro Leu Phe Ala Phe>

      3030      3040      3050      3060      3070
      *         *         *         *         *
GAA TGG AGA GAT CCA GGT ACG GGA AGA ACC GGG CAG CTC ACC TGG ACC
Glu Trp Arg Asp Pro Gly Thr Gly Arg Thr Gly Gln Leu Thr Trp Thr>

```

FIGURE 3,CONT.

26/34

(SEQ ID NO: 3)
cont'd

```

      3080      3090      3100      3110
      *      *      *      *
CGA CTG CCC CAA GGG TTC AAG AAC TCC CCG ACC ATC TTT GAC GAA GCC
Arg Leu Pro Gln Gly Phe Lys Asn Ser Pro Thr Ile Phe Asp Glu Ala>

3120      3130      3140      3150      3160
      *      *      *      *      *
CTA CAC AGA GAC CTG GCC AAC TTC AGG ATC CAA CAC CCT CAG GTG ACC
Leu His Arg Asp Leu Ala Asn Phe Arg Ile Gln His Pro Gln Val Thr>

3170      3180      3190      3200      3210
      *      *      *      *      *
CTC CTC CAG TAC GTG GAT GAC CTG CTT CTG GCG GGA GCC ACC AAA CAG
Leu Leu Gln Tyr Val Asp Asp Leu Leu Leu Ala Gly Ala Thr Lys Gln>

      3220      3230      3240      3250      3260
      *      *      *      *      *
GAC TGC TTA GAA GGC ACG AAG GCA CTA CTG CTG GAA TTG TCT GAC CTA
Asp Cys Leu Glu Gly Thr Lys Ala Leu Leu Leu Glu Leu Ser Asp Leu>

      3270      3280      3290      3300      3310
      *      *      *      *      *
GCC TAC AGA GCC TCT GCT AAG AAG GCC CAG ATT TGC AGG AGA GAG GTA
Gly Tyr Arg Ala Ser Ala Lys Lys Ala Gln Ile Cys Arg Arg Glu Val>

      3320      3330      3340      3350
      *      *      *      *
ACA TAC TTG GCG TAC AGT TTG CCG GAC GCG CAG CGA TGG CTG ACG GAG
Thr Tyr Leu Gly Tyr Ser Leu Arg Asp Gly Gln Arg Trp Leu Thr Glu>

3360      3370      3380      3390      3400
      *      *      *      *      *
GCA CCG AAG AAA ACT GTA GTC CAG ATA CCG GCC CCA ACC ACA GCC AAA
Ala Arg Lys Lys Thr Val Val Gln Ile Pro Ala Pro Thr Thr Ala Lys>

3410      3420      3430      3440      3450
      *      *      *      *      *
CAA ATG AGA GAG TTT TTG GGG ACA GCT GGA TTT TGC AGA CTG TGG ATC
Gln Met Arg Glu Phe Leu Gly Thr Ala Gly Phe Cys Arg Leu Trp Ile>

      3460      3470      3480      3490      3500
      *      *      *      *      *
CCG GCG TTT CCG ACC TTA GCA GCC CCA CTC TAC CCG CTA ACC AAA GAA
Pro Gly Phe Ala Thr Leu Ala Ala Pro Leu Tyr Pro Leu Thr Lys Glu>

      3510      3520      3530      3540      3550
      *      *      *      *      *
AAA GGG GAA TTC TCC TGG GCT CCT GAG CAC CAG AAG GCA TTT GAT GCT
Lys Gly Glu Phe Ser Trp Ala Pro Glu His Gln Lys Ala Phe Asp Ala>

      3560      3570      3580      3590
      *      *      *      *
ATC AAA AAG GCC CTG CTG AGC CCA CCT GCT CTG GCC CTC CCT GAC GTA
Ile Lys Lys Ala Leu Leu Ser Ala Pro Ala Leu Ala Leu Pro Asp Val>

3600      3610      3620      3630      3640
      *      *      *      *      *
ACT AAA CCC TTT ACC CTT TAT GTG GAT GAG CGT AAG GGA GTA GCC CCG
Thr Lys Pro Phe Thr Leu Tyr Val Asp Glu Arg Lys Gly Val Ala Arg>

```

FIGURE 3,CONT.

27/34

(SEQ ID NO: 3)
cont'd

```

3650      3660      3670      3680      3690
*          *          *          *          *
GGA GTT TTA ACC CAA ACC CTA GGA CCA TGG AGA AGA OCT GTC GCC TAC
Gly Val Leu Thr Gln Thr Leu Gly Pro Trp Arg Arg Pro Val Ala Tyr>

3700      3710      3720      3730      3740
*          *          *          *          *
CTG TCA AAG AAG CTC GAT CCT GTA GCC AGT GGT TGG CCC ATA TGC CTG
Leu Ser Lys Lys Leu Asp Pro Val Ala Ser Gly Trp Pro Ile Cys Leu>

3750      3760      3770      3780      3790
*          *          *          *          *
AAG GCT ATC GCA GCT GTG GCC ATA CTG GTC AAG GAC GCT GAC AAA TTG
Lys Ala Ile Ala Ala Val Ala Ile Leu Val Lys Asp Ala Asp Lys Leu>

3800      3810      3820      3830
*          *          *          *
ACT TTG GGA CAG AAT ATA ACT GTA ATA GCC CCC CAT GCA TTG GAG AAC
Thr Leu Gly Gln Asn Ile Thr Val Ile Ala Pro His Ala Leu Glu Asn>

3840      3850      3860      3870      3880
*          *          *          *          *
ATC GTT CCG CAG CCC CCA GAC CGA TGG ATG ACC AAC GCC CGC ATG ACC
Ile Val Arg Gln Pro Pro Asp Arg Trp Met Thr Asn Ala Arg Met Thr>

3890      3900      3910      3920      3930
*          *          *          *          *
CAC TAT CAA AGC CTG CTT CTC ACA GAG AAG GTC ACG TTC GCT CCA CCA
His Tyr Gln Ser Leu Leu Leu Thr Glu Arg Val Thr Phe Ala Pro Pro>

3940      3950      3960      3970      3980
*          *          *          *          *
GCC GCT CTC AAC CCT GCC ACT CTT CTG CCT GAA GAG ACT GAT GAA CCA
Ala Ala Leu Asn Pro Ala Thr Leu Leu Pro Glu Glu Thr Asp Glu Pro>

3990      4000      4010      4020      4030
*          *          *          *          *
GTG ACT CAT GAT TGC CAT CAA CTA TTG ATT GAG GAG ACT CGG GTC CGC
Val Thr His Asp Cys His Gln Leu Leu Ile Glu Glu Thr Gly Val Arg>

4040      4050      4060      4070
*          *          *          *
AAG GAC CTT ACA GAC ATA CCG CTG ACT GGA GAA GTG CTA ACC TGG TTC
Lys Asp Leu Thr Asp Ile Pro Leu Thr Gly Glu Val Leu Thr Trp Phe>

4080      4090      4100      4110      4120
*          *          *          *          *
ACT GAC GGA AGC AGC TAT GTG GTG GAA GGT AAG AGG ATG GCT GGG GCG
Thr Asp Gly Ser Ser Tyr Val Val Glu Gly Lys Arg Met Ala Gly Ala>

4130      4140      4150      4160      4170
*          *          *          *          *
GGG GTG GTG GAC GGG ACC CGC ACG ATC TGG GCG AGC AGC CTG CCG GAA
Ala Val Val Asp Gly Thr Arg Thr Ile Trp Ala Ser Ser Leu Pro Glu>

4180      4190      4200      4210      4220
*          *          *          *          *
GGA ACT TCA GCA CAA AAG GCT GAG CTC ATG GCC CTC ACG CAA GCT TTG
Gly Thr Ser Ala Gln Lys Ala Glu Leu Met Ala Leu Thr Gln Ala Leu>

```

FIGURE 3,CONT.

28/34

(SEQ ID NO: 3)
cont'd

```

      4230      4240      4250      4260      4270
      *        *        *        *        *
CGG CTG GCC GAA GGG AAA TCC ATA AAC ATT TAT ACG GAC AGC AGG TAT
Arg Leu Ala Glu Gly Lys Ser Ile Asn Ile Tyr Thr Asp Ser Arg Tyr>

      4280      4290      4300      4310
      *        *        *        *
GCC TTT GCG ACT GCA CAC GTA CAT GGG GCC ATC TAT AAA CAA AGG GGG
Ala Phe Ala Thr Ala His Val His Gly Ala Ile Tyr Lys Gln Arg Gly>

4320      4330      4340      4350      4360
*        *        *        *        *
TTG CTT ACC TCA GCA GGG AGG GAA ATA AAG AAC AAA GAG GAA ATT CTA
Leu Leu Thr Ser Ala Gly Arg Glu Ile Lys Asn Lys Glu Glu Ile Leu>

4370      4380      4390      4400      4410
*        *        *        *        *
AGC CTA TTA GAA GCC GTA CAT TTA CCA AAA AGG CTA GCT ATT ATA CAC
Ser Leu Leu Glu Ala Val His Leu Pro Lys Arg Leu Ala Ile Ile His>

      4420      4430      4440      4450      4460
      *        *        *        *        *
TGT CCT GGA CAT CAG AAA GCT AAA GAT CTC ATA TCC AGA GGA AAC CAG
Cys Pro Gly His Gln Lys Ala Lys Asp Leu Ile Ser Arg Gly Asn Gln>

      4470      4480      4490      4500      4510
      *        *        *        *        *
ATG GCT GAC CGG GTT CCC AAG CAG GCA GCC CAC CTT GGT AAC CTT CTG
Met Ala Asp Arg Val Ala Lys Gln Ala Ala Gln Gly Val Asn Leu Leu>

      4520      4530      4540      4550
      *        *        *        *
CCT ATA ATA GAA ATG CCC AAA GCC CCA GAA CCC AGA CGA CAG TAC ACC
Pro Ile Ile Glu Met Pro Lys Ala Pro Glu Pro Arg Arg Gln Tyr Thr>

4560      4570      4580      4590      4600
*        *        *        *        *
CTA GAA GAC TGG CAA GAG ATA AAA AAG ATA GAC CAG TTC TCT GAG ACT
Leu Glu Asp Trp Gln Glu Ile Lys Lys Ile Asp Gln Phe Ser Glu Thr>

4610      4620      4630      4640      4650
*        *        *        *        *
CCG GAA GGG ACC TCC TAT ACC TCA GAT GGG AAG GAA ATC CTG CCC CAC
Pro Glu Gly Thr Cys Tyr Thr Ser Asp Gly Lys Glu Ile Leu Pro His>

      4660      4670      4680      4690      4700
      *        *        *        *        *
AAA GAA GGG TTA GAA TAT GTC CAA CAG ATA CAT CGT CTA ACC CAC CTA
Lys Glu Gly Leu Glu Tyr Val Gln Gln Ile His Arg Leu Thr His Leu>

      4710      4720      4730      4740      4750
      *        *        *        *        *
CGA ACT AAA CAC CTG CAG CAG TTG GTC AGA ACA TCC CCT TAT CAT GTT
Gly Thr Lys His Leu Gln Gln Leu Val Arg Thr Ser Pro Tyr His Val>

      4760      4770      4780      4790
      *        *        *        *
CTG AGG CTA CCA GGA GTG GCT GAC TCG CTG GTC AAA CAT TGT GTG CCC
Leu Arg Leu Pro Gly Val Ala Asp Ser Val Val Lys His Cys Val Pro>

```

FIGURE 3,CONT.

29/34

(SEQ ID NO: 3)
cont'd

```

4800      4810      4820      4830      4840
*         *         *         *         *
TGC CAG CTG GTT AAT CCT AAT CCT TCC AGA ATG CCT CCA GGG AAG AGA
Cys Gln Leu Val Asn Ala Asn Pro Ser Arg Met Pro Pro Gly Lys Arg>

4850      4860      4870      4880      4890
*         *         *         *         *
CTA AGG GGA ACC CAC CCA GGC GCT CAC TGG GAA GTG GAC TTC ACT GAG
Leu Arg Gly Ser His Pro Gly Ala His Trp Glu Val Asp Phe Thr Glu>

4900      4910      4920      4930      4940
*         *         *         *         *
GTA AAG CCG GCT AAA TAC GGA AAC AAA TAC CTA TTG GTT TTT GTA GAC
Val Lys Pro Ala Lys Tyr Gly Asn Lys Tyr Leu Leu Val Phe Val Asp>

4950      4960      4970      4980      4990
*         *         *         *         *
ACC TTT TCA GGA TGG GTA GAG GCT TAT CCT ACT AAG AAA GAG ACT TCA
Thr Phe Ser Gly Trp Val Glu Ala Tyr Pro Thr Lys Lys Glu Thr Ser>

5000      5010      5020      5030
*         *         *         *
ACC GTG GTG GCT AAA AAA ATA CTG GAA GAA ATT TTT CCA AGA TTT GGA
Thr Val Val Ala Lys Lys Ile Leu Glu Glu Ile Phe Pro Arg Phe Gly>

5040      5050      5060      5070      5080
*         *         *         *         *
ATA CCT AAC CTT ATA GGG TCA GAC AAT GGT CCA GCT TTT GTT GCC CAG
Ile Pro Lys Val Ile Gly Ser Asp Asn Gly Pro Ala Phe Val Ala Gln>

5090      5100      5110      5120      5130
*         *         *         *         *
GTA AGT CAG GGA CTG GCC AAG ATA TTG GGG ATT GAT TGG AAA CTG CAT
Val Ser Gln Gly Leu Ala Lys Ile Leu Gly Ile Asp Trp Lys Leu His>

5140      5150      5160      5170      5180
*         *         *         *         *
TGT GCA TAC AGA CCC CAA AGC TCA GGA CAG GTA GAG AGG ATG AAT AGA
Cys Ala Tyr Arg Pro Gln Ser Ser Gly Gln Val Glu Arg Met Asn Arg>

5190      5200      5210      5220      5230
*         *         *         *         *
ACC ATT AAA GAG ACC CTT ACT AAA TTG ACC GCG GAG ACT GCC GTT AAT
Thr Ile Lys Glu Thr Leu Thr Lys Leu Thr Ala Glu Thr Gly Val Asn>

5240      5250      5260      5270
*         *         *         *
GAT TGG ATA CCT CTC CTG CCC TTT GTG CTT TTT AGG GTT AGG AAC ACC
Asp Trp Ile Ala Leu Leu Pro Phe Val Leu Phe Arg Val Arg Asn Thr>

5280      5290      5300      5310      5320
*         *         *         *         *
CCT GGA CAG TTT GGG CTG ACC CCC TAT GAA TTA CTC TAC GGG GGA CCC
Pro Gly Gln Phe Gly Leu Thr Pro Tyr Glu Leu Leu Tyr Gly Gly Pro>

5330      5340      5350      5360      5370
*         *         *         *         *
CCC CCA TTG GTA GAA ATT GCT TCT GTA CAT AGT GCT GAC GTG CTG CTT
Pro Pro Leu Val Glu Ile Ala Ser Val His Ser Ala Asp Val Leu Leu>

```

FIGURE 3,CONT.

30/34

(SEQ ID NO: 3)
cont'd

```

5380      5390      5400      5410      5420
*         *         *         *         *
TCC CAG CCT TTG TTC TCT ACG CTC AAG GCA CTT GAG TGG GTG AGA CAA
Ser Gln Pro Leu Phe Ser Arg Leu Lys Ala Leu Glu Trp Val Arg Gln>

      5430      5440      5450      5460      5470
*         *         *         *         *
CGA GCG TGG AGG CAA CTC CGG GAG GCC TAC TCA GGA GGA GGA GAC TTG
Arg Ala Trp Arg Gln Leu Arg Glu Ala Tyr Ser Gly Gly Gly Asp Leu>

      5480      5490      5500      5510
*         *         *         *         *
CAG ATC CCA CAT CGT TTC CAA GTG GGA GAT TCA GTC TAC GTT AGA CGC
Gln Ile Pro His Arg Phe Gln Val Gly Asp Ser Val Tyr Val Arg Arg>

5520      5530      5540      5550      5560
*         *         *         *         *
CAC CGT GCA GGA AAC CTC GAG ACT CCG TGG AAG GGC CCT TAT CTC GTA
His Arg Ala Gly Asn Leu Glu Thr Arg Trp Lys Gly Pro Tyr Leu Val>

5570      5580      5590      5600      5610
*         *         *         *         *
CTT TTG ACC ACA CCA ACG GCT GTG AAA GTC GAA GGA ATC TCC ACC TGG
Leu Leu Thr Thr Pro Thr Ala Val Lys Val Glu Gly Ile Ser Thr Trp>

      5620      5630      5640      5650      5660
*         *         *         *         *
ATC CAT GCA TCC CAC GTT AAA CCG GCG CCA CCT CCC GAT TCG GGG TGG
Met His Pro Thr Leu Asn Arg Arg His Leu Pro Ile Arg Gly Gly>
Ile His Ala Ser His Val Lys Pro Ala Pro Pro Pro Asp Ser Gly Trp>

      5670      5680      5690      5700      5710
*         *         *         *         *
AAA GCC GAA AAG ACT GAA AAT CCC CTT AAG CTT CGC CTC CAT CGC GTG
Lys Pro Lys Arg Leu Lys Ile Pro Leu Ser Phe Ala Ser Ile Ala Trp>
Lys Ala Glu Lys Thr Glu Asn Pro Leu Lys Leu Arg Leu His Arg Val>

      5720      5730      5740      5750      5760
*         *         *         *         *
GTT CCT TAC TCT GTC AAT AAC CTC TCA GAC T AAT GGT ATG CGC ATA GGA
Phe Leu Thr Leu Ser Ile Thr Ser Gln Thr Asn Gly Met Arg Ile Gly>
Val Pro Tyr Ser Val Asn Asn Leu Ser Asp>

      5770      5780      5790      5800
*         *         *         *
GAC AGC CTG AAC TCC CAT AAA CCC TTA TCT CTC ACC TGG TTA ATT ACT
Asp Ser Leu Asn Ser His Lys Pro Leu Ser Leu Thr Trp Leu Ile Thr>

5810      5820      5830      5840      5850
*         *         *         *         *
GAC TCC GGC ACA GGT ATT AAT ATC AAC AAC ACT CAA GGG GAG GCT CCT
Asp Ser Gly Thr Gly Ile Asn Ile Asn Asn Thr Gln Gly Glu Ala Pro>

```

FIGURE 3,CONT.

31/34

(SEQ ID NO: 3)
cont'd

```

5860      5870      5880      5890      5900
*         *         *         *         *
TTA GGA ACC TGG TGG CCT GAT CTA TAC GTT TGC CTC AGA TCA GTT ATT
Leu Gly Thr Trp Trp Pro Asp Leu Tyr Val Cys Leu Arg Ser Val Ile>

5910      5920      5930      5940      5950
*         *         *         *         *
CCT AGT CTG ACC TCA CCC CCA GAT ATC CTC CAT GCT CAC GGA TTT TAT
Pro Ser Leu Thr Ser Pro Pro Asp Ile Leu His Ala His Gly Phe Tyr>

5960      5970      5980      5990      6000
*         *         *         *         *
GTT TGC CCA GGA CCA CCA AAT AAT GGA AAA CAT TGC GGA AAT CCC AGA
Val Cys Pro Gly Pro Pro Asn Asn Gly Lys His Cys Gly Asn Pro Arg>

6010      6020      6030      6040
*         *         *         *
GAT TTC TTT TGT AAA CAA TGG AAC TGT GTA ACC TCT AAT GAT GGA TAT
Asp Phe Phe Cys Lys Gln Trp Asn Cys Val Thr Ser Asn Asp Gly Tyr>

6050      6060      6070      6080      6090
*         *         *         *         *
TGG AAA TGG CCA ACC TCT CAG CAG GAT AGG GTA AGT TTT TCT TAT GTC
Trp Lys Trp Pro Thr Ser Gln Gln Asp Arg Val Ser Phe Ser Tyr Val>

6100      6110      6120      6130      6140
*         *         *         *         *
AAC ACC TAT ACC AGC TCT GGA CAA TTT AAT TAC CTG ACC TGG ATT AGA
Asn Thr Tyr Thr Ser Ser Gly Gln Phe Asn Tyr Leu Thr Trp Ile Arg>

6150      6160      6170      6180      6190
*         *         *         *         *
ACT GGA AGC CCC AAG TGC TCT CCT TCA GAC CTA GAT TAC CTA AAA ATA
Thr Gly Ser Pro Lys Cys Ser Pro Ser Asp Leu Asp Tyr Leu Lys Ile>

6200      6210      6220      6230      6240
*         *         *         *         *
AGT TTC ACT GAG AAA GGA AAA CAA GAA AAT ATC CTA AAA TGG GTA AAT
Ser Phe Thr Glu Lys Gly Lys Gln Glu Asn Ile Leu Lys Trp Val Asn>

6250      6260      6270      6280
*         *         *         *
GGT ATG TCT TGG GGA ATG GTA TAT TAT GGA GGC TCG GGT AAA CAA CCA
Gly Met Ser Trp Gly Met Val Tyr Tyr Gly Gly Ser Gly Lys Gln Pro>

6290      6300      6310      6320      6330
*         *         *         *         *
GGC TCC ATT CTA ACT ATT CGC CTC AAA ATA AAC CAG CTG GAG CCT CCA
Gly Ser Ile Leu Thr Ile Arg Leu Lys Ile Asn Gln Leu Glu Pro Pro>

6340      6350      6360      6370      6380
*         *         *         *         *
ATG GCT ATA GGA CCA AAT ACG GTC TTG ACG GGT CAA AGA CCC CCA ACC
Met Ala Ile Gly Pro Asn Thr Val Leu Thr Gly Gln Arg Pro Pro Thr>

6390      6400      6410      6420      6430
*         *         *         *         *
CAA GGA CCA GGA CCA TCC TCT AAC ATA ACT TCT GGA TCA GAC CCC ACT
Gln Gly Pro Gly Pro Ser Ser Asn Ile Thr Ser Gly Ser Asp Pro Thr>

```

FIGURE 3.CONT.

32/34

(SEQ ID NO: 3)
cont'd

```

      644C      6450      6460      6470      6480
      *         *         *         *         *
GAG TCT AAC AGC ACG ACT AAA ATG GGG GCA AAA CTT TTT AGC CTC ATC
Glu Ser Asn Ser Thr Thr Lys Met Gly Ala Lys Leu Phe Ser Leu Ile>

      6490      6500      6510      6520
      *         *         *         *
CAG GGA CCT TTT CAA GCT CTT AAC TCC ACC ACT CCA GAG GCT ACC TCT
Gln Gly Ala Phe Gln Ala Leu Asn Ser Thr Thr Pro Glu Ala Thr Ser>

6530      6540      6550      6560      6570
      *         *         *         *         *
TCT TGT TGG CTA TGC TTA GCT TCG GGC CCA CCT TAC TAT GAA GGA ATG
Ser Cys Trp Leu Cys Leu Ala Ser Gly Pro Pro Tyr Tyr Glu Gly Met>

      6580      6590      6600      6610      6620
      *         *         *         *         *
GCT AGA AGA GGG AAA TTC AAT GTG ACA AAA GAA CAT AGA GAC CAA TGC
Ala Arg Arg Gly Lys Phe Asn Val Thr Lys Glu His Arg Asp Gln Cys>

      6630      6640      6650      6660      6670
      *         *         *         *         *
ACA TGG GGA TCC CAA AAT AAG CTT ACC CTT ACT GAG GTT TCT GGA AAA
Thr Trp Gly Ser Gln Asn Lys Leu Thr Leu Thr Glu Val Ser Gly Lys>

      6680      6690      6700      6710      6720
      *         *         *         *         *
GGC ACC TGC ATA GGA AAG GTT CCC CCA ACC CAC CAA CAC CTT TGT AAC
Gly Thr Cys Ile Gly Lys Val Pro Pro Ser His Gln His Leu Cys Asn>

      6730      6740      6750      6760
      *         *         *         *
CAC ACT GAA GCC TTT AAT CAA ACC TCT GAG AGT CAA TAT CTG GTA CCT
His Thr Glu Ala Phe Asn Gln Thr Ser Glu Ser Gln Tyr Leu Val Pro>

6770      6780      6790      6800      6810
      *         *         *         *         *
GGT TAT GAC AGG TGG TGG GCA TGT AAT ACT GGA TTA ACC CCT TGT GTT
Gly Tyr Asp Arg Trp Trp Ala Cys Asn Thr Gly Leu Thr Pro Cys Val>

      6820      6830      6840      6850      6860
      *         *         *         *         *
TCC ACC TTG GTT TTT AAC CAA ACT AAA GAT TTT TGC ATT ATG GTC CAA
Ser Thr Leu Val Phe Asn Gln Thr Lys Asp Phe Cys Ile Met Val Gln>

      6870      6880      6890      6900      6910
      *         *         *         *         *
ATT GTT CCC CGA GTG TAT TAC TAT CCC GAA AAA CCA ATC CTT GAT GAA
Ile Val Pro Arg Val Tyr Tyr Tyr Pro Glu Lys Ala Ile Leu Asp Glu>

      6920      6930      6940      6950      6960
      *         *         *         *         *
TAT GAC TAC AGA AAT CAT CGA CAA AAG AGA GAA CCC ATA TCT CTG ACA
Tyr Asp Tyr Arg Asn His Arg Gln Lys Arg Glu Pro Ile Ser Leu Thr>

      6970      6980      6990      7000
      *         *         *         *
CTT GCT GTG ATG CTC GGA CTT GGA GTG GCA GCA GGT GTA GGA ACA GGA
Leu Ala Val Met Leu Gly Leu Gly Val Ala Ala Gly Val Gly Thr Gly>

```

FIGURE 3,CONT.

33/34

(SEQ ID NO: 3)
cont'd

7010 7020 7030 7040 7050
* * * * *
ACA GCT GCC CTG GTC ACG GGA CCA CAG CAG CTA GAA ACA GGA CTT AGT
Thr Ala Ala Leu Val Thr Gly Pro Gln Gln Leu Glu Thr Gly Leu Ser>
7060 7070 7080 7090 7100
* * * * *
AAC CTA CAT CGA ATT GTA ACA GAA GAT CTC CAA GCC CTA GAA AAA TCT
Asn Leu His Arg Ile Val Thr Glu Asp Leu Gln Ala Leu Glu Lys Ser>
7110 7120 7130 7140 7150
* * * * *
GTC AGT AAC CTG GAG GAA TCC CTA ACC TCC TTA TCT GAA GTA GTC CTA
Val Ser Asn Leu Glu Glu Ser Leu Thr Ser Leu Ser Glu Val Val Leu>
7160 7170 7180 7190 7200
* * * * *
CAG AAT AGA AGA GGG TTA GAT TTA TTA TTT CTA AAA GAA GGA GGA TTA
Gln Asn Arg Arg Gly Leu Asp Leu Leu Phe Leu Lys Glu Gly Gly Leu>
7210 7220 7230 7240
* * * *
TGT GTA GCC TGG AAG GAG GAA TGC TGT TTT TAT GTG GAT CAT TCA GCG
Cys Val Ala Leu Lys Glu Glu Cys Cys Phe Tyr Val Asp His Ser Gly>
7250 7260 7270 7280 7290
* * * * *
GCC ATC ACA GAC TCC ATG AAC AAG CTT AGA GAA AGG TTG GAG AAG CGT
Ala Ile Arg Asp Ser Met Asn Lys Leu Arg Glu Arg Leu Glu Lys Arg>
7300 7310 7320 7330 7340
* * * * *
CGA AGG GAA AAG GAA ACT ACT CAA GGG TGG TTT CAG GGA TGG TTC AAC
Arg Arg Glu Lys Glu Thr Thr Gln Gly Trp Phe Glu Gly Trp Phe Asn>
7350 7360 7370 7380 7390
* * * * *
AGG TCT CTT TGG TTG GCT ACC CTA CTT TCT GCT TTA ACA GGA CCC TTA
Arg Ser Leu Trp Leu Ala Thr Leu Leu Ser Ala Leu Thr Gly Pro Leu>
7400 7410 7420 7430 7440
* * * * *
ATA GTC CTC CTC CTG TTA CTC ACA GTT GGG CCA TGT ATT ATT AAC AAG
Ile Val Leu Leu Leu Leu Leu Thr Val Gly Pro Cys Ile Ile Asn Lys>
7450 7460 7470 7480
* * * *
TTA ATT GCC TTC ATT AGA GAA CGA ATA AGT GCA GTC CAG ATC ATG GTA
Leu Ile Ala Phe Ile Arg Glu Arg Ile Ser Ala Val Gln Ile Met Val>
7490 7500 7510 7520 7530
* * * * *
CTT AGA CAA CAG TAC CAA ACC CCG TCT AGC AGG GAA GCT GGC CCG
Leu Arg Gln Gln Tyr Gln Ser Pro Ser Ser Arg Glu Ala Gly Arg>
7540 7550 7560 7570 7580 7590
* * * * * *
TAGCTCT ACCAGTTCTA AGATTAGAAC TATTACAAAG AGAAGAAGTG GGGAAATGAAA

FIGURE 3.CONT.

34/34

(SEQ ID NO: 3)
cont'd

```

      7600      7610      7620      7630      7640      7650
      * *      * *      * *      * *      * *      * *
GGATGAAAAT ACAACCTAAG CTAATGAGAA GCTTAAAAAT GTTCTGAATT CCAGAGTTTG

      7660      7670      7680      7690      7700      7710
      * *      * *      * *      * *      * *      * *
TTCTTATAG GTAAAAGATT AGGTTTTTTG CTGTTTAAAT ATATGCOGAA GTAAAATAGG

      7720      7730      7740      7750      7760      7770
      * *      * *      * *      * *      * *      * *
CCCTGAGTAC ATGTCTCTAG GCATGAAACT TCTTGAAACT ATTTGAGATA ACAAGAAAAG

      7780      7790      7800      7810      7820      7830
      * *      * *      * *      * *      * *      * *
GGAGTTTCTA ACTGCTTGTT TAGCTTCTGT AAAACTGGTT GCGCCATAAA GATGTTGAAA

      7840      7850      7860      7870      7880      7890
      * *      * *      * *      * *      * *      * *
TGTGTGATACA CATATCTTGG TGACAACATG TCTCCCCCAC CCCGAAACAT GCGCAAATGT

      7900      7910      7920      7930      7940      7950
      * *      * *      * *      * *      * *      * *
GTAACICTAA AACAATTAA ATTAATTGGT CCACGAAGCG CGGCTCTCG AAGTTTTAAA

      7960      7970      7980      7990      8000      8010
      * *      * *      * *      * *      * *      * *
TTGACTGGTT TGTGATATTT TGAAATGATT GGTTTGTAAA GCGCGGCTT TGTCTGAAC

      8020      8030      8040      8050      8060      8070
      * *      * *      * *      * *      * *      * *
CCCATAAAAG CTGTCCCGAC TCACACTCG GGGCCGAGT CCTCTACCCG TGGGTGGTGT

      8080      8090      8100      8110      8120      8130
      * *      * *      * *      * *      * *      * *
ACGACTGTGG GCCCAGGCC GCTTGAATA AAAATCCTCT TCTGTTTGTC ATCAAAAAAA

AA

```

FIGURE 3,CONT.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/19680

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
1-14, 28, and 38
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/19630

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12Q 1/68, 1/70

US CL : 435/5, 6; 536/22.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/5, 6; 536/22.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, STN

search terms: swine, retrovirus, DNA, nucleic acid, hybridization

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|----------------------------|
| X | BOWES. Localization of a retroviral element within the rd gene coding the Beta Subunit of cGMP phosphodiesterase Proc. Natl. Acad. Sci. USA. April 1993. Vol. 90, pages 2955-2959, especially page 2958.1, 4-9, 11, 12, 28, and 38 | 1, 4-9, 11, 12, 28, and 38 |
| X | DELASSUS et al. Genetic Organization of Gibbon Ape Leukemia Virus, Virology. 1989. Vol. 173, pages 205-213, especially pages 207-208. | 1, 4-9, 11, 12, 28, and 38 |



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z

document member of the same patent family

Date of the actual completion of the international search

03 APRIL 1997

Date of mailing of the international search report

30 APR 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JEZIA RILEY

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/19680

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|------------------------------|
| X | DEVARE et al. Nucleotide Sequence of the Simian Sarcoma Virus Genome: Demonstration that its Acquired Cellular Sequences Encode the Transforming Gene Product p28 Proc. Natl. Acad. Sci. USA. February 1983. Vol 80. pages 731-735, especially page 732. | 1, 4-9, 11-14, 28, and 38 |